

# Washington Flora Checklist

## A checklist of the Vascular Plants of Washington State

### Hosted by the University of Washington Herbarium

#### Family: Asteraceae

462 terminal taxa (species, subspecies, and varieties).

The Washington Flora Checklist aims to be a complete list of the native and naturalized vascular plants of Washington State, with current classifications, nomenclature and synonymy.

#### Taxa included in the checklist:

- \* Native taxa whether extant, extirpated, or extinct.
- \* Exotic taxa that are naturalized, escaped from cultivation, or persisting wild.
- \* Waifs (e.g., ballast plants, escaped crop plants) and other scarcely collected exotics.
- \* Interspecific hybrids that are frequent or self-maintaining.
- \* Some unnamed taxa in the process of being described.

Family classifications follow [APG IV](#) for angiosperms, PPG I (J. Syst. Evol. 54:563?603. 2016.) for pteridophytes, and Christenhusz et al. (Phytotaxa 19:55?70. 2011.) for gymnosperms, with a few exceptions. Nomenclature and synonymy at the rank of genus and below follows the [2nd Edition of the Flora of the Pacific Northwest](#) except where superceded by new information.

Accepted names are indicated with blue font; synonyms with black font.  
Native species and infraspecies are marked with **boldface** font.

**Please note:** This is a working checklist, continuously updated. Use it at your discretion.

Created from the Washington Flora Checklist Database on May 19th, 2024 at 12:39pm PST.  
Available online at <https://burkeherbarium.org/waflora/>

Comments and questions should be addressed to the checklist administrators:  
David Giblin ([dgiblin@uw.edu](mailto:dgiblin@uw.edu))  
Peter Zika ([zikap941@gmail.com](mailto:zikap941@gmail.com))

#### Suggested citation:

Weinmann, F., P.F. Zika, D.E. Giblin, B. Legler. 2002+. Checklist of the Vascular Plants of Washington State. University of Washington Herbarium. <https://burkeherbarium.org/waflora/>. Accessed 19 May 2024.

## Dicots:

### Asteraceae [FNA19, HC2] Aster Family

#### Synonyms:

Compositae [HC]

Many taxonomic and nomenclatural changes have occurred within Asteraceae since the publication of Hitchcock and Cronquist (1973). The Flora of North America project(FNA) published the Asteraceae volumes in 2007, and that has served as the primary literature resource for the taxonomy and nomenclature provided here. Some of the introduced taxa in Washington belonging to this family are not included in the FNA volumes. Many of these can be found in Stace's New Flora of the British Isles (1997). Ken Chambers and Scott Sundberg provided a treatment of the Asteraceae for the Oregon Flora Project (OFP), and some of their taxonomic decisions are followed here rather than what is provided in FNA.

#### References:

(none)

#### *Achillea* [FNA19, HC, HC2]

Sp. Pl. 2: 896. 1753; Gen. Pl. ed. 5, 382. 1754.

yarrow

#### *Achillea millefolium* L. [FNA19, HC, HC2]

Sp. Pl. 2: 899.

milfoil, yarrow

*Achillea borealis* Bong.

*Achillea lanulosa* Nutt. var. *eradiata* (Piper) M. Peck

*Achillea lanulosa* Nutt. var. *lanulosa*

*Achillea millefolium* L. ssp. *lanulosa* (Nutt.) Piper [HC]

*Achillea millefolium* L. var. *alpicola* (Rydb.) Garrett [HC]

*Achillea millefolium* L. var. *borealis* (Bong.) Farw.

*Achillea millefolium* L. var. *californica* (Pollard) Jeps. [HC]

*Achillea millefolium* L. var. *lanulosa* (Nutt.) Piper [HC]

*Achillea millefolium* L. var. *litoralis* Ehrendorfer ex Nobs

*Achillea millefolium* L. var. *millefolium*

*Achillea millefolium* L. var. *occidentalis* DC.

*Achillea millefolium* L. var. *pacifica* (Rydb.) G.N. Jones

#### *Achillea ptarmica* L. [FNA19, HC2]

Sp. Pl. 2: 898.

pearl yarrow

FNA19 lists this species as occurring in WA. No voucher, reported by R. Old in Kz99. FNA19: "Achillea ptarmica is naturalized from Eurasia. "Double-flowered" plants originated as cultivars; apparently, they persist outside of cultivation."

#### *Adenocaulon* [FNA19, HC, HC2]

Bot. Misc. 1: 19, plate 15. 1829.

pathfinder, trail plant

#### *Adenocaulon bicolor* Hook. [FNA19, HC, HC2]

Bot. Misc. 1: 19, plate 15.

pathfinder, trailplant

#### *Ageratina* [FNA21, HC2]

Hist. Nat. Vég. 10: 286. 1841.

snakeroot

#### *Ageratina occidentalis* (Hook.) R.M. King & H. Rob. [FNA21, HC2]

Phytologia. 19: 224.

western boneset, western snakeroot

*Eupatorium occidentale* Hook. [HC]

***Agoseris* [FNA19, HC, HC2]**

Fl. Ludov. 58. 1817.

false-dandelion, mountain-dandelion

***Agoseris apargioides* (Less.) Greene [FNA19, HC, HC2]**

Pittonia. 2: 177.

seaside agoseris

**var. *maritima* (E. Sheldon) G.I. Baird [FNA19, HC, HC2]**

Sida. 21: 716.

seaside agoseris

*Agoseris apargioides* (Less.) Greene ssp. *maritima* (E. Sheldon) Q. Jones

*Agoseris maritima* E. Sheldon

***Agoseris aurantiaca* (Hook.) Greene [FNA19, HC, HC2]**

Pittonia. 2: 177.

orange agoseris

**var. *aurantiaca* [FNA19, HC, HC2]**

orange agoseris, slender agoseris

*Agoseris angustissima* Greene

*Agoseris arachnoidea* Rydb.

*Agoseris aurantiaca* (Hook.) Greene ssp. *aurantiaca*

*Agoseris gracilens* (A. Gray) Greene

*Agoseris greenei* (A. Gray) Rydb.

*Agoseris howellii* Greene

*Agoseris nana* Rydb.

*Agoseris prionophylla* Greene

*Agoseris subalpina* G.N. Jones

*Agoseris vulcanica* Greene

FNA19:"Variety aurantiaca is widespread in the western cordillera and is disjunct in Quebec. Two morphologic trends occur within this variety. Plants of wetter habitats represent the typical var. aurantiaca; those of drier habitats resemble what past authors have called *Agoseris gracilens* (including *A. gracilens* var. *greenei*). There is a weak geographic trend to this variation, with the aurantiaca phase occurring mostly along the Rocky Mountains axis and the gracilens phase mostly along the Cascade Mountains-Sierra Nevada axis. In their extremes they appear distinct, but their intergradation is so complete that separation becomes arbitrary. Putative hybrids between var. aurantiaca and *A. glauca*, *A. grandiflora*, *A. monticola*, and *A. parviflora* have been collected. Corolla color in var. aurantiaca is variable but most commonly orange. Pink-flowered forms occur sporadically. They have been recognized as *Agoseris lackschewitzii*. Recognition of pink forms is unmerited; if it were, the older name *A. carnea* would have priority."

**var. *carnea* P. Lesica [HC2]**

Journal of Botanical Research Institute of Texas 6(1): 25-27.

pink agoseris

*Agoseris lackschewitzii* Douglas M. Hend. & R.K. Moseley

***Agoseris x elata* (Nutt.) Greene [FNA19, HC, HC2]**

Pittonia. 2: 177.

tall agoseris, tall goat-chicory

*Agoseris laciniata* (Nutt.) Greene

***Agoseris glauca* (Pursh) Raf. [FNA19, HC, HC2]**

Herb. Raf. 39.

pale agoseris, short-beaked agoseris

(see also *Agoseris agrestis*, *Agoseris monticola*)

var. *dasycephala* (Torr. & A. Gray) Jeps. [FNA19, HC, HC2]

Man. Fl. Pl. Calif. 1005.

pale goat-chicory

*Agoseris glauca* (Pursh) Raf. var. *aspera* (Rydb.) Cronquist

FNA19: "Variety *dasycephala* occurs primarily at high elevations in the western cordillera, extending eastward onto the northern prairies, and disjunctively in the Canadian arctic (Caribou Hills). It is more readily distinguished from var. *glauca* southward, where the two varieties are ± elevationally separated. Difficulty in separating them occurs northward, where they are nearer each other and pockets of complete introgression occur, e.g., southeastern British Columbia and southwestern Alberta. Hybrids with *Agoseris aurantiaca* and *A. parviflora* also occur. Variety *dasycephala* contains regional phases that exhibit a step-clinal distribution. The large number of synonyms reflects the variation. As circumscribed here, var. *dasycephala* encompasses most of what has been called *Agoseris glauca* var. *agrestis* (see discussion under var. *glauca*)."

var. *glauca* [FNA19, HC, HC2]

Herb. Raf. 39.

*Agoseris xagrestis* Osterh. [HC2]

*Agoseris lacera* Greene

*Agoseris lopathifolia* Greene

*Agoseris longissima* Greene

*Agoseris microdonta* Greene

*Agoseris procera* Greene

*Agoseris vicinalis* Greene

FNA19: "Variety *glauca* is usually found at lower elevations from the northern prairies westward to valleys and basins of the North American cordillera. Misidentification is often due to falsely assuming this variety is strictly glabrous. Some regional phases have a high percentage of individuals with weakly puberulent peduncles and/or phyllaries. In addition, var. *glauca* intergrades with var. *dasycephala* in some locations."

***Agoseris grandiflora* (Nutt.) Greene [FNA19, HC, HC2]**

Pittonia. 2: 178.

large-flowered agoseris

*Stylopappus grandiflorus* Nutt.

var. *grandiflora* [FNA19, HC2]

Pittonia. 2: 178.

large flowered agoseris, large flower goat-chicory

*Agoseris cinerea* Greene

*Agoseris grandiflora* (Nutt.) Greene var. *intermedia* (Greene) Jeps.

*Agoseris grandiflora* (Nutt.) Greene var. *plebeia* (Greene) G.L. Wittrock

*Agoseris intermedia* Greene

*Agoseris marshallii* (Greene) Greene

*Agoseris obtusifolia* (Suksd.) Rydb.

*Agoseris plebeia* (Greene) Greene

FNA19: "Variety *grandiflora* is most commonly found east of the Cascade Mountains and southward into California and occurs primarily in grassland, steppe, or chaparral. It has regional phases, especially southward in its range. These appear more or less distinct but they so completely intergrade that their separation becomes arbitrary. Variety *grandiflora* rarely forms intermediates with other species; putative hybrids with *A. apargioides* have been collected. It is one of the suspected parental taxa of *A. xelata*, especially the Sierra Nevada populations."

var. *leptophylla* G.I. Baird [FNA19, HC2]

Sida. 21: 267.

Puget Sound agoseris

FNA19: "Variety *leptophylla* is most commonly found west of the Cascade Mountains from Vancouver Island through the Puget Sound and Willamette Valley to the Siskiyou-Klamath Mountains region of southwestern Oregon and northwestern California. It also occurs sporadically in mesic forest areas on

the eastern slopes of the Cascade Mountains, and disjunctively in the Selkirk-Clearwater Mountains region of British Columbia and northern Idaho. In the Selkirk-Clearwater Mountains region, Columbia River Gorge, southern Willamette Valley, and Siskiyou-Klamath Mountains region var. *grandiflora* and var. *leptophylla* are sympatric and appear to be introgressive. In those regions, intermediate specimens are not uncommon. It may be one of the parental taxa of *A. xelata* (which see), especially the Puget Sound-Willamette Valley populations."

***Agoseris heterophylla* (Nutt.) Greene [FNA19, HC, HC2]**

Pittonia. 2: 178.

annual agoseris

*Agoseris heterophylla* (Nutt.) Greene ssp. *heterophylla*

var. *heterophylla* [FNA19, HC, HC2]

*Agoseris heterophylla* (Nutt.) Greene ssp. *normalis* Piper

***Agoseris monticola* Greene [FNA19, HC2]**

Pittonia. 4: 37.

mountain agoseris, Sierra Nevada agoseris

*Agoseris glauca* (Pursh) Raf. var. *monticola* (Greene) Q. Jones [HC]

FNA19: "Agoseris monticola occurs mainly in the Sierra Nevada and sporadically eastward in the Great Basin (Jarbridge and Ruby Mountains) and northward to the Cascade Range and Blue Mountains of Oregon. It appears to be allied with *A. glauca* and has been treated as a variety of the latter. Ecologically, it approaches *A. glauca* var. *dasycephala*; the two are morphologically and geographically separate from each other. Intermediates between *A. monticola* and *A. aurantiaca*, *A. glauca*, and *A. parviflora* are known."

***Agoseris retrorsa* (Benth.) Greene [FNA19, HC, HC2]**

Pittonia. 2: 178. 1891.

spear leaved agoseris, spear leaf goat-chicory

*Macrorhynchus angustifolius* Kellogg

*Macrorhynchus retrorsus* Benth.

***Ambrosia* [FNA21, HC, HC2]**

Sp. Pl. 2: 987. 1753; Gen. Pl. ed. 5, 425. 1754.

bursage, burweed, ragweed

***Ambrosia acanthicarpa* Hook. [FNA21, HC, HC2]**

Fl. Bor.-Amer. 1: 309.

flat spine bur-ragweed, annual bursage, bur ragweed

*Franseria acanthicarpa* (Hook.) Coville

***Ambrosia artemisiifolia* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 988.

annual ragweed, common ragweed

*Ambrosia artemisiifolia* L. var. *elatior* (L.) Descourtilz

*Ambrosia artemisiifolia* L. var. *paniculata* (Michx.) Blank.

*Ambrosia elatior* L.

*Ambrosia glandulosa* Scheele

*Ambrosia monophylla* (Walter) Rydb.

FNA21: "Hybrids between *Ambrosia psilostachya* and *A. artemisiifolia* have been called *A. xintergradiens* W. H. Wagner." FNA21: "The name *Ambrosia xhelena* Rouleau applies to hybrids between *A. artemisiifolia* and *A. trifida*."

***Ambrosia chamissonis* (Less.) Greene [FNA21, HC, HC2]**

Man. Bot. San Francisco. 188.

silver beachweed, beach bur, cutleaf beach bur, silver bur-ragweed

*Ambrosia chamissonis* (Less.) Greene var. *bipinnatisecta* (Less.) J.T. Howell [HC]

*Ambrosia chamissonis* (Less.) Greene var. *chamissonis* [HC]

*Franseria chamissonis* Less. ssp. *bipinnatisecta* (Less.) Wiggins & Stockw.  
*Franseria chamissonis* Less. ssp. *chamissonis*  
*Franseria chamissonis* Less. var. *bipinnatisecta* Less.  
*Franseria chamissonis* Less. var. *chamissonis*

***Ambrosia psilostachya* DC. [FNA21, HC, HC2]**

Prodr. 5: 526.  
perennial ragweed, western ragweed

*Ambrosia psilostachya* DC. var. *californica* (Rydb.) S.F. Blake  
*Ambrosia psilostachya* DC. var. *coronopifolia* (Torr. & A. Gray) Farw.  
*Ambrosia psilostachya* DC. var. *lindheimeriana* (Scheele) Blank.  
*Ambrosia rugelii* Rydb.

FNA21: "Hybrids between *Ambrosia psilostachya* and *A. artemisiifolia* have been called *A. ×intergradiens* W. H. Wagner."

***Ambrosia trifida* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 987.  
giant ragweed

*Ambrosia aptera* DC.  
*Ambrosia trifida* L. var. *integrifolia* (Muhl. ex Willd.) Torr. & A. Gray  
*Ambrosia trifida* L. var. *texana* Scheele  
*Ambrosia trifida* L. var. *trifida* [HC]

***Anaphalis* [FNA19, HC, HC2]**

Prodr. 6: 271. 1838.  
pearly-everlasting

***Anaphalis margaritacea* (L.) Benth. & Hook. f. [FNA19, HC, HC2]**

Gen. Pl. 2: 303.  
pearly everlasting

*Anaphalis margaritacea* (L.) Benth. & Hook. f. var. *occidentalis* Greene  
*Anaphalis margaritacea* (L.) Benth. & Hook. f. var. *subalpina* (A. Gray) A. Gray  
*Gnaphalium margaritaceum* L.

***Anisocarpus* [FNA21, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 388. 1841.  
anisocarpus

***Anisocarpus madioides* Nutt. [FNA21, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 388.  
woodland tarplant, woodland tarweed  
*Madia madioides* (Nutt.) Greene [HC]

***Antennaria* [FNA19, HC, HC2]**

Fruct. Sem. Pl. 2: 410, plate 167, fig. 3. 1791.  
everlasting, pussy-toes

***Antennaria alpina* (L.) Gaertn. [FNA19, HC, HC2]**

Fruct. Sem. Pl. 2: 410.  
alpine pussytoes

*Antennaria alpina* (L.) Gaertn. var. *canescens* Lange  
*Gnaphalium alpinum* L.

The application of this species concept to plants in Washington warrants further consideration. . FNA19: "Excluded names: Some *Antennaria* names are based on early-generation interspecific hybrids, including: *A. ×rousseauui* A. E. Porsild = ? *A. alpina* × *A. rosea* *Antennaria alpina* is one of the more morphologically variable agamic complexes in the genus. Some taxonomists have argued that true *Antennaria alpina* does not occur in North America, because none of the North American material exactly matches the type of *A. alpina*, which is from Lapland (M. O. Malte 1934; A. E. Porsild 1965). If one uses a strict typological

species concept, then this is true; I recognize that this species complex is composed of innumerable apomictic clones and am circumscribing a broad species concept for *A. alpina*. The potential morphologic overlap between the *A. media* and *A. alpina* complexes is a major taxonomic problem. The chief difference between members of the two complexes is the presence of prominent flags on cauline leaves in *A. alpina* and their absence in *A. media*. *Antennaria alpina* of North America is gynoecious and characterized by its dark green to black phyllaries and conspicuous flags on the distal cauline leaves. The basal leaves vary from glabrous, as in the type material, to pubescent. The primary progenitors of the *A. alpina* complex include *A. aromatica*, *A. densifolia*, *A. friesiana* subsp. *alaskana*, *A. friesiana* subsp. *neoalaskana*, *A. monocephala* subsp. *monocephala*, and *A. pulchella*."

***Antennaria anaphaloides* Rydb. [FNA19, HC, HC2]**

Mem. New York Bot. Gard. 1: 409.

tall pussytoes

*Antennaria anaphaloides* Rydb. var. *straminea* B. Boivin

*Antennaria pulcherrima* (Hook.) Greene ssp. *anaphaloides* (Rydb.) W.A. Weber

*Antennaria pulcherrima* (Hook.) Greene var. *anaphaloides* (Rydb.) G.W. Douglas

***Antennaria corymbosa* E.E. Nelson [FNA19, HC, HC2]**

Bot. Gaz. 27: 212. 1899.

flat topped pussytoes, meadow pussytoes

*Antennaria acuta* Rydb.

*Antennaria dioica* (L.) Gaertn. var. *corymbosa* (E.E. Nelson) Jeps.

*Antennaria hygrophila* Greene

*Antennaria nardina* Greene

Scarcely different from *A. rosea*. FNA19: "Antennaria corymbosa is characterized by linear-ob lanceolate basal leaves and white-tipped phyllaries, each with a distinct black spot near the base of the scarious portion. A form with black phyllaries (*A. acuta*) occurs sporadically throughout the range of the species (R. J. Bayer 1988). *Antennaria corymbosa* is a sexual progenitor of the *A. rosea* complex."

***Antennaria dimorpha* (Nutt.) Torr. & A. Gray [FNA19, HC, HC2]**

Fl. N. Amer. 2: 431.

cushion pussytoes, low pussytoes

*Antennaria dimorpha* (Nutt.) Torr. & A. Gray var. *integra* L.F. Hend.

*Antennaria dimorpha* (Nutt.) Torr. & A. Gray var. *latisquama* (Piper) M. Peck

*Antennaria dimorpha* (Nutt.) Torr. & A. Gray var. *macrocephala* D.C. Eaton

*Antennaria dimorpha* (Nutt.) Torr. & A. Gray var. *nuttallii* D.C. Eaton

*Antennaria latisquama* Piper

*Antennaria macrocephala* (D.C. Eaton) Rydb.

*Gnaphalium dimorphum* Nutt.

***Antennaria flagellaris* (A. Gray) A. Gray [FNA19, HC, HC2]**

Proc. Amer. Acad. Arts. 17: 212.

stoloniferous pussytoes, whip pussytoes

*Antennaria dimorpha* (Nutt.) Torr. & A. Gray var. *flagellaris* A. Gray

***Antennaria geyeri* A. Gray [FNA19, HC, HC2]**

Mem. Amer. Acad. Arts, n. s. 4: 107.

Geyer's pussytoes, pinewoods pussytoes

FNA19: "Antennaria geyeri is distinctive because it has woody upright branches and is not stoloniferous. It lacks basal leaves at flowering and has heads that are often described as subdioecious (central flowers are often bisexual). As the only member of the Geyerae group, *A. geyeri* is not closely related to any other species of *Antennaria*; it bears strong similarities to some species of *Anaphalis* (R. J. Bayer 1990; Bayer et al. 1996)."

***Antennaria howellii* Greene [FNA19, HC2]**

Pittonia. 3: 174.

Howell's pussytoes

**ssp. *howellii* [FNA19, HC2]**

Pittonia. 3: 174.

Howell's pussytoes

*Antennaria neglecta* Greene ssp. *howellii* (Greene) Hultén

*Antennaria neglecta* Greene var. *howellii* (Greene) Cronquist [HC]

*Antennaria neodioica* Greene ssp. *howellii* (Greene) Bayer

ssp. *neodioica* (Greene) R.J. Bayer [FNA19, HC2]

Brittonia. 41: 397.

northern pussytoes

*Antennaria howellii* Greene ssp. *petaloidea* (Fernald) R.J. Bayer [FNA19]

*Antennaria neglecta* Greene var. *attenuata* (Fernald) Cronquist [HC]

*Antennaria neglecta* Greene var. *neodioica* (Greene) Cronquist

*Antennaria pedicellata* Greene

***Antennaria lanata* (Hook.) Greene [FNA19, HC, HC2]**

Pittonia. 3: 288.

woolly pussytoes

*Antennaria carpatica* (Wahlenb.) Hook. var. *lanata* Hook., orthographic variant

***Antennaria luzuloides* Torr. & A. Gray [FNA19, HC, HC2]**

Fl. N. Amer. 2: 430.

silvery-brown pussytoes, woodrush pussytoes

ssp. *luzuloides* [FNA19, HC2]

silvery brown pussytoes, woodrush pussytoes

*Antennaria argentea* Benth. ssp. *argentea*

*Antennaria luzuloides* Torr. & A. Gray var. *luzuloides*

***Antennaria media* Greene [FNA19, HC2]**

Pittonia. 3: 286.

alpine pussytoes, Rocky Mountain pussytoes

*Antennaria alpina* (L.) Gaertn. var. *media* (Greene) Jeps. [HC]

FNA19: "The main distinction between *A. media* and *A. alpina* is flags on distal caudine leaves present in *A. alpina* and mostly absent in *A. media* (Bayer 1990d). Phyllaries of the pistillate plants in *A. alpina* tend to be acute; they are blunter in *A. media*. At some point, it may be preferable to follow W. L. Jepson ([1923?1925]) and some later authors and treat *A. media* as a subspecies of *A. alpina*. *Antennaria media* appears to be an autopolyploid derivative of *A. pulchella*; genes from *A. pulchella* may have introgressed into the *A. alpina* and *A. parvifolia* complexes indirectly through *A. media*."

***Antennaria microphylla* Rydb. [FNA19, HC, HC2]**

Bull. Torrey Bot. Club. 24: 303.

little-leaf pussytoes, rosy pussytoes, white pussytoes

*Antennaria bracteosa* Rydb.

*Antennaria concinna* E.E. Nelson

*Antennaria microphylla* Lunell var. *soltstitialis* Lunell

*Antennaria nitida* Greene

*Antennaria rosea* Greene [FNA19]

*Antennaria rosea* Greene ssp. *arida* (E.E. Nelson) R.J. Bayer [FNA19]

*Antennaria rosea* Greene ssp. *confinis* (Greene) R.J. Bayer [FNA19]

*Antennaria rosea* Greene ssp. *pulvinata* (Greene) R.J. Bayer [FNA19]

*Antennaria rosea* Greene ssp. *rosea* [FNA19]

*Antennaria rosea* Greene var. *nitida* (Greene) Breitung

*Antennaria solstitialis* Lunell

FNA19: "Antennaria microphylla is a primary sexual progenitor of the *A. rosea* polyploid agamic complex (R. J. Bayer 1990b). A. Cronquist (1955) included *A. rosea* within his circumscription of *A. microphylla*. It is preferable to recognize sexual diploids as distinct from their morphologically discrete hybrid apomictic derivatives. *Antennaria microphylla* is always dioecious and has stems distally stipitate-glandular and white phyllaries; *A. rosea* is always gynoecious and has stems without glandular hairs and phyllaries only occasionally white. Some authors (A. E. Porsild 1950; E. H. Moss 1959; Porsild and W. J. Cody 1980)

have recognized *A. nitida* as distinct; comparisons of the nomenclatural types of the two show that they are conspecific. *Antennaria microphylla* has allelopathic properties (G. D. Manners and D. S. Galitz 1985)."

***Antennaria monocephala* DC. [FNA19, HC2]**

Prodri. 6: 269.

pygmy pussytoes, single-headed pussytoes

Recently (2017) photographed in Glacier Peak Wilderness. Identification confirmed by Jamie Fenneman at UBC.

***Antennaria parvifolia* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 406.

little-leaf pussytoes, Nuttall's pussytoes

***Antennaria pulcherrima* (Hook.) Greene [FNA19, HC, HC2]**

Pittonia. 3: 176.

showy pussytoes

*Antennaria pulcherrima* (Hook.) Greene ssp. *eucosma* (Fernald & Wiegand) R.J. Bayer [FNA19]

*Antennaria pulcherrima* (Hook.) Greene ssp. *pulcherrima* [FNA19]

*Antennaria pulcherrima* (Hook.) Greene var. *pulcherrima*

***Antennaria pulvinata* Greene**

Pittonia 3: 287.

white pussytoes

***Antennaria racemosa* Hook. [FNA19, HC, HC2]**

Fl. Bor.-Amer. 1: 330.

Hooker's pussytoes, raceme pussytoes

***Antennaria stenophylla* (A. Gray) A. Gray [FNA19, HC, HC2]**

Proc. Amer. Acad. Arts. 17: 213.

narrowleaved pussytoes

*Antennaria alpina* (L.) Gaertn. var. *stenophylla* A. Gray

*Antennaria leucophaea* Piper

***Antennaria umbrinella* Rydb. [FNA19, HC, HC2]**

Bull. Torrey Bot. Club. 24: 302.

brown-bract pussytoes, umber pussytoes

***Anthemis* [FNA19, HC, HC2]**

Sp. Pl. 2: 893. 1753; Gen. Pl. ed. 5, 381. 1754.

chamomile, dogfennel, mayweed

(see also *Cota*)

***Anthemis arvensis* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 894.

corn chamomile, field chamomile

*Anthemis arvensis* L. var. *arvensis*

***Anthemis cotula* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 894.

mayweed chamomile, stinking chamomile, dogfennel

***Arctium* [FNA19, HC, HC2]**

Sp. Pl. 2: 816. 1753; Gen. Pl. ed. 5, 357. 1754.

burdock, clotbur

***Arctium lappa* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 816.

great burdock, greater burdock

***Arctium minus* (Hill) Bernh. [FNA19, HC, HC2]**

Syst. Verz. 154.

common burdock, lesser burdock

***Arnica*** [FNA21, HC, HC2]

Sp. Pl. 2: 884. 1753; Gen. Pl. ed. 5, 376. 1754.  
arnica

***Arnica chamissonis*** Less. [FNA21, HC, HC2]

Linnaea. 6: 238.  
leafy arnica, meadow arnica, narrowleaf arnica, silvery arnica, leafy leapordbane  
*Arnica chamissonis* Less. ssp. *chamissonis* [HC]  
*Arnica chamissonis* Less. ssp. *foliosa* (Nutt.) Maguire [HC]  
*Arnica chamissonis* Less. ssp. *incana* (A. Gray) Maguire  
*Arnica chamissonis* Less. var. *andina* (Nutt.) Ediger & T.M. Barkl.  
*Arnica chamissonis* Less. var. *chamissonis*  
*Arnica chamissonis* Less. var. *foliosa* (Nutt.) Maguire [HC]  
*Arnica chamissonis* Less. var. *incana* (A. Gray) Hultén [HC]  
*Arnica chamissonis* Less. var. *interior* Maguire [HC]  
*Arnica chamissonis* Less. var. *maguirei* (A. Nels.) Maguire [HC]

***Arnica cordifolia*** Hook. [FNA21, HC, HC2]

Fl. Bor.-Amer. 1: 331.  
heart-leaf arnica, heart-leaf leopardbane  
*Arnica cordifolia* Hook. var. *cordifolia* [HC]  
*Arnica cordifolia* Hook. var. *pumila* (Rydb.) Maguire [HC]

***Arnica discoidea*** Benth. [FNA21, HC]

Pl. Hartw. 319.  
rayless arnica, rayless leopardbane  
*Arnica discoidea* Benth. var. *eradiata* (A. Gray) Cronquist [HC]  
*Arnica grayi* A. Heller  
*Arnica parviflora* A. Gray ssp. *alata* (Rydb.) Maguire  
*Arnica parviflora* A. Gray ssp. *parviflora*

***Arnica fulgens*** Pursh [FNA21, HC, HC2]

Fl. Amer. Sept. 2: 527.  
hillside arnica, orange arnica, shining leopardbane

***Arnica gracilis*** Rydb. [FNA21, HC2]

Bull. Torrey Bot. Club. 24: 297.  
slender arnica, slender leopardbane  
*Arnica latifolia* Bong. var. *gracilis* (Rydb.) Cronquist [HC]

***Arnica lanceolata*** Nutt. [FNA21, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 407.  
clasping arnica, stream bank arnica  
ssp. *prima* (Maguire) Strother & S.J. Wolf [FNA21, HC2]  
Fl. N. Amer. (1993+). 21: 373.  
clasping arnica, streambank arnica, streambank leopardbane  
*Arnica amplexicaulis* Nutt. [HC]  
*Arnica amplexicaulis* Nutt. ssp. *amplexicaulis*  
*Arnica amplexicaulis* Nutt. var. *amplexicaulis* [HC]  
*Arnica amplexicaulis* Nutt. var. *piperi* H. St. John & F.A. Warren [HC]  
*Arnica amplexifolia* Rydb. ssp. *prima* Maguire

***Arnica latifolia*** Bong. [FNA21, HC, HC2]

Mém. Acad. Imp. Sci. St.-Pétersbourg, Sér. 6, Sci. Math. 2: 147.  
broad-leaved arnica, mountain arnica, daffodil leopardbane  
(see also *Arnica gracilis*)

*Arnica latifolia* Bong. var. *latifolia* [HC]

***Arnica longifolia*** D.C. Eaton [FNA21, HC, HC2]

Botany (Fortieth Parallel). 186.

longleaf arnica, seep spring arnica, spear-leaf leopardbane

*Arnica longifolia* D.C. Eaton ssp. *myriadenia* (Piper) Maguire

***Arnica mollis*** Hook. [FNA21, HC, HC2]

Fl. Bor.-Amer. 1: 331.

cordilleran arnica, hairy arnica, cordilleran leopardbane

***Arnica nevadensis*** A. Gray [FNA21, HC, HC2]

Proc. Amer. Acad. Arts. 19: 55.

Nevada arnica, Sierra arnica, Sierran leopardbane

*Arnica tomentella* Greene

***Arnica ovata*** Greene [FNA21, HC2]

Pittonia. 4: 161.

sticky arnica, sticky-leaf arnica

*Arnica ×diversifolia* Greene [HC]

***Arnica parryi*** A. Gray [FNA21, HC, HC2]

Amer. Naturalist. 8: 213.

Parry's arnica

*Arnica angustifolia* Vahl ssp. *eradiata* A. Gray

*Arnica parryi* A. Gray ssp. *parryi*

*Arnica parryi* A. Gray ssp. *sonnei* (Greene) Maguire

*Arnica parryi* A. Gray var. *parryi* [HC]

*Arnica parryi* A. Gray var. *sonnei* (Greene) Cronquist

***Arnica rydbergii*** Greene [FNA21, HC, HC2]

Pittonia. 4: 36.

Rydberg's arnica, subalpine arnica, subalpine leopardbane

***Arnica sororia*** Greene [FNA21, HC, HC2]

Ottawa Naturalist. 23: 213.

bunch arnica, twin arnica, twin leopardbane

*Arnica fulgens* Pursh var. *sororia* (Greene) G.W. Douglas & Ruyle-Douglas

***Artemisia*** [FNA19, HC, HC2]

Sp. Pl. 2: 845. 1753; Gen. Pl. ed. 5, 367. 1754.

artemisia, mugwort, sagebrush, wormwood

*Picrothamnus* [FNA19]

*Sphaeromeria* [FNA19]

***Artemisia absinthium*** L. [FNA19, HC, HC2]

Sp. Pl. 2: 848.

absinthe, old-man, wormwood

*Artemisia absinthium* L. var. *absinthium*

***Artemisia annua*** L. [FNA19, HC, HC2]

Sp. Pl. 2: 847.

sweet Annie, sweet sagewort, annual wormwood

***Artemisia arbuscula*** Nutt. [FNA19, HC, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 398.

dwarf sagebrush, low sagebrush

ssp. *arbuscula* [FNA19, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 398.

little sagebrush, low sagebrush

*Artemisia arbuscula* Nutt. var. *arbuscula* [HC]

FNA19:"The relatively large heads of *Artemisia arbuscula* subsp. *arbuscula* suggest a relationship with *A. cana*; the extreme morphologic variability of this subspecies from east to west may be the result of

hybridization with various subspecies within the *A. cana* complex."

***Artemisia biennis* Willd. [FNA19, HC, HC2]**

Phytographia. 11.  
biennial wormwood

*Artemisia biennis* Willd. var. *biennis*

FNA19: "Artemisia biennis is naturalized and weedy in the eastern portion of its range. It is morphologically similar to *A. annua*, differing primarily in the coarser leaf lobes and larger heads that are sessile in axils of leaflike bracts. *Artemisia biennis* is considered native to the northwest United States; it may be introduced in other parts of its range. The type specimen is a horticultural specimen from New Zealand."

***Artemisia campestris* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 846.  
Pacific sagewort, northern wormwood

var. *borealis* (Pall.) M. Peck [HC2]

Man. Pl. Oregon 768.  
northern wormwood

*Artemisia borealis* Pall. [FNA19]

*Artemisia borealis* Pall. ssp. *borealis* [FNA19]

*Artemisia borealis* Pall. ssp. *richardsoniana* (Besser) Korobkov [FNA19]

*Artemisia campestris* L. ssp. *borealis* (Pall.) H.M. Hall & Clem. [HC]

*Artemisia campestris* L. var. *purshii* (Besser) Cronquist [HC]

var. *scouleriana* (Besser) Cronquist [HC, HC2]

Leaflets of Western Botany 7(2): 20.  
Pacific sagewort, Scouler's wormwood

*Artemisia campestris* L. ssp. *pacifica* (Nutt.) H.M. Hall & Clem. [FNA19]

var. *wormskioldei* (Besser ex Hook.) Cronquist [HC, HC2]

Leafl. W. Bot. 6: 43.  
Columbia Islands sagewort, Wormskiolde's wormwood sagewort

***Artemisia cana* Pursh [FNA19, HC, HC2]**

Fl. Amer. Sept. 2: 521.  
hoary sagebrush, silver sagebrush

ssp. *bolanderi* (A. Gray) G.H. Ward [FNA19, HC2]

Contr. Dudley Herb. 4: 192.  
Bolander's hairy sagebrush

***Artemisia douglasiana* Besser [FNA19, HC, HC2]**

Fl. Bor.-Amer. 1: 323.  
Douglas's mugwort, Douglas's sagewort, Douglas's wormwood

*Artemisia vulgaris* L. var. *douglasiana* (Besser) H. St. John

***Artemisia dracunculus* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 849.  
dragon sagewort, tarragon, dragon wormwood

*Artemisia dracunculus* L. ssp. *dracunculus*

*Artemisia dracunculus* L. var. *dracunculus* [HC]

***Artemisia frigida* Willd. [FNA19, HC, HC2]**

Sp. Pl. 3: 1838.  
prairie sagebrush, prairie sagewort

***Artemisia furcata* M. Bieb. [FNA19, HC2]**

Fl. Taur.-Caucas. 3: 567.  
three-forked mugwort, forked wormwood

*Artemisia furcata* M. Bieb. var. *furcata*

*Artemisia furcata* M. Bieb. var. *heterophylla* (Besser) Hultén

*Artemisia trifurcata* Stephani ex Spreng. [HC]

FNA19: "Artemisia furcata extends from the islands of the Bering Sea into southern and interior Alaska, parts of Canada (disjunct in British Columbia and the northernmost Rocky Mountains of Alberta), and on Mt. Rainier in Washington. The array of names applied to A. furcata shows the taxonomic confusion arising from a myriad of morphologic variants that may indicate introgression with other species."

***Artemisia ludoviciana* Nutt. [FNA19, HC, HC2]**

Gen. N. Amer. Pl. 2: 143.

western mugwort, prairie sage

**ssp. *candicans* (Rydb.) D.D. Keck [FNA19, HC2]**

Proc. Calif. Acad. Sci., ser. 4. 25: 447.

gray sagewort

*Artemisia ludoviciana* Nutt. var. *latiloba* Nutt. [HC]

**ssp. *incompta* (Nutt.) D.D. Keck [FNA19, HC2]**

Publ. Carnegie Inst. Wash. 520: 327.

intermediate sagewort, mountain wormwood

*Artemisia ludoviciana* Nutt. var. *incompta* (Nutt.) Cronquist [HC]

**ssp. *lindleyana* (Besser) Lesica [HC2]**

J. Bot. Res. Inst. Texas 6(1): 26

Lindley's western mugwort, Lindley's prairie sage

*Artemisia lindleyana* Besser. [HC]

**ssp. *ludoviciana* [FNA19, HC2]**

Nouv. Mém. Soc. Imp. Naturalistes Moscou 3: 38.

western mugwort, Louisiana sagewort, silver wormwood

*Artemisia diversifolia* Rydb.

*Artemisia gnaphalooides* Nutt.

*Artemisia ludoviciana* Nutt. var. *ludoviciana* [HC]

***Artemisia michauxiana* Besser [FNA19, HC, HC2]**

Fl. Bor.-Amer. 1: 324.

Michaux's mugwort, lemon sagewort, Michaux's wormwood

*Artemisia vulgaris* L. var. *michauxiana* (Besser) H. St. John

FNA19: "Members of the Artemisia ludoviciana complex with deeply lobed leaves are sometimes confused with A. michauxiana, and there is evidence that plants hybridize in some locations. Artemisia michauxiana is distinguished by its glabrous, bright green to yellow-green foliage and leemony-sweet fragrance."

***Artemisia norvegica* Fr. [FNA19, HC, HC2]**

Novit. Fl. Svec. 56. 1817.

mountain sagewort, boreal wormwood

**ssp. *saxatilis* (Besser) H.M. Hall & Clem. [FNA19, HC2]**

Publ. Carnegie Inst. Wash. 326: 58. 1923.

mountain sagewort

*Artemisia arctica* Less. ssp. *arctica*

*Artemisia norvegica* Fr. var. *saxatilis* (Besser) Jeps. [HC]

*Artemisia saxatalis* Less.

***Artemisia rigida* (Nutt.) A. Gray [FNA19, HC, HC2]**

Proc. Amer. Acad. Arts. 19: 49.

scabland sagebrush, stiff sagebrush

***Artemisia spiciformis* Osterh. [FNA19, HC2]**

Bull. Torrey Bot. Club. 27: 507.

snowfield sagebrush, spiked sagebrush

*Artemisia tridentata* Nutt. ssp. *spiciformis* (Osterh.) Kartesz & Gandhi

FNA19: "Often confused with *Artemisia rothrockii*, *A. spiciformis* has been recognized only recently as a widespread, high-elevation sagebrush of late-lying snowfields. Molecular analysis has not yet determined the degree to which this species intergrades with *A. cana* subsp. *viscidula* and *A. tridentata* subsp. *vaseyana*, the presumed parents of this putative hybrid. Because snow-field sagebrush produces fertile seeds and forms a stable community type, it is treated here as a distinct species."

***Artemisia stelleriana* Besser [FNA19, HC2]**

Nouv. Mém. Soc. Imp. Naturalistes Moscou. 3: 79, plate 5.  
dusty miller, old-woman, beach wormwood, Steller's wormwood

***Artemisia suksdorffii* Piper [FNA19, HC, HC2]**

Bull. Torrey Bot. Club. 28: 42.  
coastal mugwort, Suksdorf's sagewort, coastal wormwood

*Artemisia vulgaris* L. var. *littoralis* Suksd.

***Artemisia tilesii* Ledeb. [FNA19, HC, HC2]**

Mém. Acad. Imp. Sci. St. Pétersbourg Hist. Acad. 5: 568.  
Aleutian mugwort, Cascade wormwood

*Artemisia elatior* (Torr. & A. Gray) Rydb.

*Artemisia hookeriana* Besser

*Artemisia tilesii* Ledeb. ssp. *elatior* (Torr. & A. Gray) Hultén

*Artemisia tilesii* Ledeb. ssp. *unaliaschcensis* (Besser) Hultén

*Artemisia tilesii* Ledeb. var. *elatior* Torr. & A. Gray

*Artemisia tilesii* Ledeb. var. *unaliaschcensis* Besser [HC]

FNA19: "*Artemisia tilesii* has a bewildering array of variation in leaf and inflorescence morphology that has been separated into four infraspecific taxa recognized in some floras. I am unable to separate these taxa consistently and am including them within a broad circumscription of the species."

***Artemisia tridentata* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 398.  
big sagebrush

**ssp. *tridentata* [FNA19, HC2]**

big sagebrush

*Artemisia tridentata* Nutt. var. *tridentata*

**ssp. *vaseyana* (Rydb.) Beetle [FNA19, HC2]**

Rhodora. 61: 83.  
mountain big sagebrush, Vasey sagebrush

*Artemisia tridentata* Nutt. var. *vaseyana* (Rydb.) B. Boivin

**ssp. *wyomingensis* Beetle & A.M. Young [FNA19, HC2]**

Rhodora. 67: 405.  
Wyoming sagebrush

*Artemisia tridentata* Nutt. var. *wyomingensis* (Beetle & A.M. Young) S.L. Welsh

*Seriphidium tridentatum* (Nutt.) W.A. Weber ssp. *wyomingense* (Beetle & A.M. Young) W.A. Weber

FNA19: "Subspecies *wyomingensis* is the common sagebrush of rocky or fine-grained soils from valleys to high plateaus in the Great Basin. It is an allotetraploid that may be derived from the populations of subsp. *tridentata* with which it occurs. Identification is based primarily on the shorter leaves of subsp. *wyomingensis*, its usually shorter stature, and its shorter flowering branches that are retained from year to year. Wyoming sagebrush may be increasing in abundance in response to increased grazing pressure and drought in the high valleys of the Great Basin."

***Artemisia tripartita* Rydb. [FNA19, HC, HC2]**

Mem. New York Bot. Gard. 1: 432.  
cut-leaf sagebrush, threetip sagebrush

**ssp. *tripartita* [FNA19, HC2]**

cutleaf sagebrush, threetip sagebrush

***Artemisia vulgaris* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 848.

mugwort, lobed wormwood

*Artemisia vulgaris* L. var. *selengensis* (Turcz. ex Besser) Maxim.

*Artemisia vulgaris* L. var. *vulgaris*

***Askellia* [HC2]**

elegant hawksbeard

***Askellia pygmaea* (Ledeb.) Sennikov [HC2]**

Komarovia 5(2): 86.

low hawksbeard

*Crepis nana* Richardson [FNA19, HC]

*Crepis nana* Richardson ssp. *nana* [HC]

*Crepis nana* Richardson ssp. *ramosa* Babc. [HC]

*Crepis nana* Richardson var. *lyratifolia* (Turcz.) Hultén

*Crepis nana* Richardson var. *ramosa* (Babc.) Cronquist

FNA19: "Crepis nana occurs in North America and northern Asia. It is recognized by the tufted, cespitose habit, elongate roots and rhizomes, and occurrence in alpine habitats. In the typical form, the plants are tufted, the stems are not leafy, and the heads are borne among the leaves. Taller specimens with elongated, leafy branches and heads borne well beyond the basal leaves are sometimes recognized as subsp. *ramosa*; these characteristics appear to be part of the normal range of variation for the species. *Crepis nana* is closely related to *C. elegans*, differing mainly in the shape of the cypselae. The cypselae of *C. nana* are almost always more columnar, wider at bases, and with broader ribs, than those of *C. elegans*."

***Baccharis* [FNA20, HC, HC2]**

Sp. Pl. 2: 860. 1753; Gen. Pl. ed. 5, 370. 1754.

baccharis

***Baccharis pilularis* DC. [FNA20, HC, HC2]**

Prodri. 5: 407.

chaparral broom, coyote brush

**ssp. *consanguinea* (DC.) C.B. Wolf [FNA20, HC2]**

Occas. Pap. Rancho Santa Ana Bot. Gard. 1: 21.

chaparral broom

*Baccharis pilularis* DC. var. *consanguinea* (DC.) Kuntze [HC]

***Balsamorhiza* [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 349. 1840.

balsamroot

***Balsamorhiza ×bonseri* H. St. John [HC2]**

hybrid balsamroot

(= *Balsamorhiza rosea* × *Balsamorhiza sagittata*)

***Balsamorhiza careyana* A. Gray [FNA21, HC, HC2]**

Mem. Amer. Acad. Arts, n. s. 4: 81.

Carey's balsamroot

*Balsamorhiza careyana* A. Gray var. *careyana* [HC]

*Balsamorhiza careyana* A. Gray var. *intermedia* Cronquist [HC]

***Balsamorhiza careyana* A. Gray × *Balsamorhiza hookeri* Nutt.**

hybrid balsamroot

***Balsamorhiza deltoidea* Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 351.

deltoid balsamroot, Puget balsamroot

***Balsamorhiza hookeri* Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 349.

hairy balsamroot, hare's head balsamroot, Hooker's balsamroot

*Balsamorhiza hirsuta* Nutt. [HC]

*Balsamorhiza hirsuta* Nutt. var. *lagocephala* W.M. Sharp

*Balsamorhiza hookeri* Nutt. var. *hirsuta* (Nutt.) A. Nelson

*Balsamorhiza hookeri* Nutt. var. *hookeri* [HC]

*Balsamorhiza hookeri* Nutt. var. *lagocephala* (W.M. Sharp) Cronquist [HC]

FNA19: "At one time or another, most species of subg. *Balsamorhiza* have been synonymized under *B. hookeri*. Nevertheless, a number of taxa are justifiably segregated as species by their morphologic differences and geographic restrictions. One might logically choose either of two taxonomies: recognizing only two species in the entire genus, one representing subg. *Arthrorhiza* and the other subg. *Balsamorhiza*, or recognizing each slightly differing population as a species. Either course results in an unsatisfactory classification. The present classification is a compromise. A knotty problem persists. A central cluster of populations from eastern Washington to southeastern California display a number of minor and locally discrete morphologies. They tend to be less isolated from each other than are the peripheral populations, although some tend to mimic the latter ones in one or more characteristics. Their evolutionary history may be involved with past hybridizations with each other or with species of subg. *Arthrorhiza*, gene drift, and polyploidy. At present, it appears impossible to reach a satisfactory classification."

*Balsamorhiza hookeri* Nutt. × *Balsamorhiza sagittata* (Pursh) Nutt.

hybrid balsamroot

*Balsamorhiza incana* Nutt. [FNA21, HC, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 350.

hoary balsamroot, woolly balsamroot

*Balsamorhiza rosea* A. Nelson & J.F. Macbr. [FNA21, HC, HC2]

Bot. Gaz. 56: 478.

rosy balsamroot

*Balsamorhiza sagittata* (Pursh) Nutt. [FNA21, HC, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 350.

arrowleaf balsamroot

*Balsamorhiza serrata* A. Nelson & J.F. Macbr. [FNA21, HC, HC2]

Bot. Gaz. 56: 479.

serrate balsamroot, toothed balsamroot

*Balsamorhiza ×terebinthacea* (Hook.) Nutt. [HC2]

wormwood balsamroot

Chambers & Sundberg use this epithet for any cross between *B. deltoidea*, *B. careyana*, and *B. sagittata*, the three deltoid-leaved taxa; see Weber (1953)

*Balsamorhiza ×tomentosa* Rydb. [HC2]

woolly hybrid balsamroot

*Bellis* [FNA20, HC, HC2]

Sp. Pl. 2: 886. 1753; Gen. Pl. ed. 5, 378. 1754.

bellis, daisy

*Bellis perennis* L. [FNA20, HC, HC2]

Sp. Pl. 2: 886.

English daisy, lawn daisy

*Bidens* [FNA21, HC, HC2]

Sp. Pl. 2: 831. 1753; Gen. Pl. ed. 5, 362. 1754.

beggar-ticks, bur-marigold, sticktight

*Bidens amplissima* Greene [FNA21, HC, HC2]

Pittonia. 4: 268.

Vancouver Island beggar-ticks

*Biden cernua* L. var. *elata* Torr. & A. Gray

*Bidens elata* (Torr. & A. Gray) Sherff

***Bidens beckii*** Torr. ex Spreng. [FNA21, HC, HC2]

Neue Entd. 2: 135.

Beck's water marigold

*Megalodonta beckii* (Torr. ex Spreng.) Greene

*Megalodonta beckii* (Torr. ex Spreng.) Greene var. *beckii*

*Megalodonta beckii* (Torr. ex Spreng.) Greene var. *hendersonii* Sherff

*Megalodonta beckii* (Torr. ex Spreng.) Greene var. *oregonensis* Sherff

***Bidens cernua*** L. [FNA21, HC, HC2]

Sp. Pl. 2: 832.

nodding beggar-ticks, bur-marigold

*Bidens cernua* L. var. *cernua*

*Bidens cernua* L. var. *elliptica* Wiegand

*Bidens cernua* L. var. *minima* (Huds.) Pursh

***Bidens connata*** Muhl. ex Willd. [FNA21, HC2]

Sp. Pl. 3: 1718.

purplestem beggars-ticks, swamp beggar-ticks

FNA21:"<i>Bidens connata</i> may be better treated as part of <i>B. tripartita</i>."

***Bidens frondosa*** L. [FNA21, HC, HC2]

Sp. Pl. 2: 832.

leafy beggar-ticks, devil's pitchfork, sticktight

***Bidens tripartita*** L. [FNA21, HC, HC2]

Sp. Pl. 2: 831.

three-lobed beggar-ticks

FNA21:"Plants with cypselae mid-nerves strongly developed (cypselae more or less strongly 4-angled and, often, tuberculate) that are treated below as *Bidens connata* have been included in *B. tripartita*, perhaps rightly so. And some botanists have included (or advocated inclusion of) *B. eatonii*, *B. heterodoxa*, and/or *B. infirma* in *B. tripartita*, as well, perhaps rightly so."

***Bidens vulgata*** Greene [FNA21, HC, HC2]

Pittonia. 4: 72.

tall beggar-ticks, western sticktight

Considered Introduced in OR & Native in BC.

***Blepharipappus*** [FNA21, HC, HC2]

Fl. Bor.-Amer. 1: 316. 1833.

eyelash tarweed

***Blepharipappus scaber*** Hook. [FNA21, HC, HC2]

Fl. Bor.-Amer. 1: 316.

blepharipappus, rough eyelashweed

*Blepharipappus scaber* Hook. ssp. *laevis* (A. Gray) D.D. Keck

*Blepharipappus scaber* Hook. ssp. *scaber*

*Blepharipappus scaber* Hook. var. *scaber* [HC]

FNA21: "<i>Blepharipappus scaber</i> is unusual among self-incompatible, continental tarweeds for occurring widely in western North America and having a relatively limited distribution in the California Floristic Province."

***Boltonia*** [FNA20, HC, HC2]

Sert. Angl. 27. 1789.

Doll's-daisy

***Boltonia asteroides*** (L.) L'Hér. [FNA20, HC, HC2]

Sert. Angl. 27.

white doll's daisy

var. *recognita* (Fernald & Griscom) Cronquist [FNA20, HC, HC2]

Bull. Torrey Bot. Club. 74: 149.  
white doll's daisy

*Boltonia latisquama* A. Gray var. *microcephala* Fernald & Griscom  
*Boltonia latisquama* A. Gray var. *occidentalis* A. Gray  
*Boltonia latisquama* A. Gray var. *recognita* Fernald & Griscom  
*Boltonia recognita* (Fernald & Griscom) G.N. Jones

***Brickellia* [FNA21, HC, HC2]**

Sketch Bot. S. Carolina. 2: 290. 1823.  
brickellbush, brickellia, thoroughwort

***Brickellia grandiflora* (Hook.) Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 287. 1840.  
tasselflower brickellbush, large flowered tasselflower, large flowered thoroughwort

*Brickellia grandiflorum* Hook.

***Brickellia microphylla* (Nutt.) A. Gray [FNA21, HC, HC2]**

Smithsonian Contr. Knowl. 3(5): 85.  
small-leaved brickellbush

**var. *microphylla* [FNA21, HC, HC2]**

Smithsonian Contr. Knowl. 3(5): 85.  
small-leaved brickellbush, small-leaved brickellia

*Brickellia microphylla* (Nutt.) A. Gray var. *watsonii* (B.L. Rob.) S.L. Welsh  
*Brickellia watsonii* B.L. Rob.

***Brickellia oblongifolia* Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 288.  
narrow-leaved brickellbush, narrow-leaved thoroughwort

**var. *oblongifolia* [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 288.  
narrowleaf brickellia

***Cacaliopsis* [FNA20, HC2]**

Proc. Amer. Acad. Arts. 19: 50. 1883.  
cacaliopsis, silvercrown

***Cacaliopsis nardosmia* (A. Gray) A. Gray [FNA20, HC2]**

Proc. Amer. Acad. Arts. 19: 50.  
silvercrown luina, tall silvercrown

*Cacalia nardosmia* A. Gray

*Cacaliopsis nardosmia* (A. Gray) A. Gray ssp. *glabrata* (Piper) Piper

*Luina nardosmia* (A. Gray) Cronquist [HC]

*Luina nardosmia* (A. Gray) Cronquist var. *glabrata* (Piper) Cronquist [HC]

***Calendula* [FNA19, HC2]**

Sp. Pl. 2: 921. 1753; Gen. Pl. ed. 5, 393. 1754.

***Canadanthus* [FNA20, HC2]**

Phytologia. 77: 250. 1995.  
mountain aster

***Canadanthus modestus* (Lindl.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 251.  
few-flowered aster, great northern aster

*Aster major* (Hook.) Porter

*Aster modestus* Lindl. [HC]

*Aster modestus* Lindl. var. *major* (Hook.) Muenscher

*Aster sayianus* Nutt.

*Aster unalaschensis* Less. ex Bong. var. *major* Hook.  
*Weberaster modestus* (Lindl.) Å. Löve & D. Löve

***Carduus* [FNA19, HC, HC2]**

Sp. Pl. 2: 820. 1753; Gen. Pl. ed. 5, 358. 1754.  
plumeless thistle

***Carduus acanthoides* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 821.  
plumeless thistle, spiny plumeless thistle  
ssp. *acanthoides* [FNA19, HC2]

***Carduus nutans* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 821.  
musk thistle, nodding thistle

***Carduus pycnocephalus* L. [FNA19, HC, HC2]**

Sp. Pl. Ed. 2., 2: 1151.  
Italian plumeless thistle

ssp. *pycnocephalus* [FNA19, HC2]

Sp. Pl. Ed. 2., 2: 1151.  
Italian plumeless thistle

***Carduus tenuiflorus* Curtis [FNA19, HC, HC2]**

Fl. Londin. 2(6,61): plate 55.  
slender flowered thistle, winged plumeless thistle

*Carduus pycnocephalus* L. var. *tenuiflorus* (Curtis) Fiori

FNA 19: "Carduus tenuiflorus has been reported from New Jersey, Texas, and Washington; I have not seen specimens from those states. *Carduus pycnocephalus* and *C. tenuiflorus* are similar annuals with small, usually tightly clustered heads. The number of heads per capitulecence is usually ultimately greater in *C. tenuiflorus*, but early season plants of this species often have only a few heads. At the end of the growing season the fruiting heads of *C. tenuiflorus* are aggregated in dense, subspheric clusters. Stem wings tend to be more pronounced in *C. tenuiflorus*. Fresh corollas of *C. pycnocephalus* are rose-purple whereas those of *C. tenuiflorus* have a more pinkish tinge, but this difference is subtle and not reliable on herbarium material. The phyllaries of *C. tenuiflorus* are membranous-margined, more or less glabrate, and lack the short, stiff, upwardly appressed trichomes of *C. pycnocephalus*. All published chromosome counts for *Carduus tenuiflorus* from both Old and New World material are the same. The two species sometimes grow in mixed populations and at times appear to intergrade. Hybridization has been reported in Europe (S. W. T. Batra et al. 1981) and is suspected to occur in California. Hybrids between *C. pycnocephalus* and *C. tenuiflorus* have been designated *Carduus xtheriotii* Rouy."

***Carthamus* [FNA19, HC, HC2]**

Sp. Pl. 2: 830. 1753; Gen. Pl. ed. 5, 361. 1754.  
distaff thistle

***Centaurea* [FNA19, HC, HC2]**

Sp. Pl. 2: 909. 1753; Gen. Pl. ed. 5, 389. 1754.  
centaurea, knapweed, star-thistle  
(see also *Rhaponticum*)

***Cnicus* [HC]**

***Centaurea benedicta* (L.) L. [FNA19, HC2]**

Sp. Pl. Ed. 2., 2: 1296.  
blessed thistle

***Cnicus benedictus* L. [HC]**

FNA19: "Recent molecular phylogenetic studies (A. Susanna et al. 1995; N. Garcia-Jacas et al. 2000, 2001) have begun to clarify relationships within *Centaurea* and between *Centaurea* and other genera. Some taxa traditionally included within *Centaurea* (e.g., the two native North American species, *Centaurea*

americana and C. rothrockii) fall outside the redefined generic boundaries and are here treated in Plectocephalus. Others usually placed into segregate genera (e.g., Cnicus benedictus) are firmly nested within Centaurea..... Although Cnicus has usually been recognized as a distinctive monotypic genus, it has been merged into Centaurea by various authors (e.g., K. Bremer 1994; G. Wagenitz and F. H. Hellwig 1996). Recent molecular systematic studies (N. Garcia-Jacas et al. 2000) provide additional evidence that it is nested within Centaurea."

*Centaurea calcitrapa* L. [FNA19, HC, HC2]

Sp. Pl. 2: 917.

purple star-thistle, red star-thistle

FNA19 includes WA within the range of this species.

*Centaurea cyanus* L. [FNA19, HC, HC2]

Sp. Pl. 2: 911.

bachelor's button, garden cornflower

*Centaurea diffusa* Lam. [FNA19, HC, HC2]

Encycl. 1: 675.

diffuse knapweed, tumble knapweed, white knapweed

*Centaurea xgerstlaueri* Erdner [FNA, HC2]

Mitt. Bayer. Bot. Ges. xxxiv. 425, ex Bot. Centralbl.xcviii. 604.

hybrid knapweed, meadow knapweed, protean knapweed

(= *Centaurea jacea* × *Centaurea nigra* or *Centaurea jacea* × *nigrescens*)

*Centaurea debeauxii* Godr. & Grenier ssp. *thuillieri* Dostál

*Centaurea xmoncktonii* C.E. Britton [FNA19]

*Centaurea nigra* L. × *Centaurea jacea* L.

*Centaurea pratensis* Thuill. [HC], superfluous renaming (illegitimate)

FNA19: "Centaurea xmoncktonii is native to Europe or originated in North America from European ancestry. Meadow knapweeds represent an array of mutually interfertile intermediates derived by hybridization and backcrossing among the various cytotypes of the *Centaurea jacea* complex. The plants variously combine features of *C. jacea* and *C. nigra*, and perhaps *C. nigrescens* as well. The hybrid complex includes both diploids and tetraploids. Extremes approach the parental types. Meadow knapweeds are often present without either parent in the immediate vicinity. They are considered to be noxious weeds in British Columbia, Idaho, Oregon, and Washington. *Centaurea pratensis* J. L. Thuillier, sometimes applied to plants that belong here, is not a legitimate name." Chambers and Sundberg (2000) treat as *C. pratensis* Thuill., which is *C. jacea* × *nigra*.

*Centaurea iberica* Trevir. ex Spreng. [FNA19, HC2]

Syst. Veg. 3: 406.

Iberian knapweed, Iberian star-thistle

FNA19: "Iberian star thistle is considered to be a noxious weed in several states of the western United States. Weed control measures in Oregon and Washington have apparently eradicated the species in those states. *Centaurea iberica* is very similar to *C. calcitrapa*, from which it differs by its pappose cypselae and often more robust habit." Chambers and Sundberg (2000) give author as Spreng. WA report is MT database Kz99, are there specimens?

*Centaurea jacea* L. [FNA19, HC, HC2]

Sp. Pl. 2: 914.

brown knapweed, brownray knapweed

*Centaurea macrocephala* Puschkarew ex Willd. [FNA19, HC2]

Sp. Pl. 3: 2298.

globe knapweed

*Centaurea melitensis* L. [FNA19, HC, HC2]

Sp. Pl. 2: 917.

Maltese star-thistle, tocalote

*Centaurea montana* L. [FNA19, HC, HC2]

Sp. Pl. 2: 911.

mountain bluet, mountain cornflower, montane star-thistle

*Centaurea nigra* L. [FNA19, HC, HC2]

Sp. Pl. 2: 911.

black knapweed, lesser knapweed

*Centaurea nigrescens* Willd. [FNA19, HC2]

Sp. Pl. 3: 2288. 1803.

short fringed knapweed, Tyrol knapweed

*Centaurea dubia* Suter [HC]

*Centaurea dubia* Suter ssp. *nigrescens* (Willd.) Hayek

*Centaurea dubia* Suter ssp. *vochinensis* (Bernh. ex Rchb.) Hayek

*Centaurea jacea* L. ssp. *nigrescens* (Willd.) Celakovsky

*Centaurea transalpina* Schleich. ex DC.

*Centaurea vochinensis* Bernh. ex Rchb.

FNA19: Tyrol knapweed is considered to be a noxious weed in Washington and Oregon. In recent years there has been much controversy regarding the name(s) to be applied to the North American Tyrol knapweeds. The names *Centaurea vochinensis*, *C. nigrescens*, and *C. dubia* have all been used in twentieth-century North American floras, and J. T. Kartesz and C. A. Meacham (1999) have accepted *C. transalpina* as well. R. J. Moore (1972) tentatively accepted two species, *C. nigrescens* and *C. dubia*, placing *C. transalpina* and *C. vochinensis* as synonyms through application beneath both species. Moore discussed the considerable similarities and practical difficulties of differentiating the taxa. H. A. Gleason and A. Cronquist (1991) recognized *C. dubia* as including *C. nigrescens* and *C. vochinensis*. E. G. Voss (1972?1996, vol. 3) recognized *C. nigrescens* as including *C. dubia* and *C. vochinensis*. Kartesz and Meacham accept *C. nigrescens* as a species, including *C. vochinensis*; they also accept *C. transalpina* with *C. dubia* as a synonym. In our investigation of the North American Tyrol knapweeds we have not been able to distinguish more than one (admittedly variable) entity. At the species level the correct name for this taxon is *Centaurea nigrescens*. *Centaurea dubia* Suter, sometimes applied to plants that belong here, is not a valid name."

*Centaurea solstitialis* L. [FNA19, HC, HC2]

Sp. Pl. 2: 917.

yellow star-thistle

*Centaurea stoebe* L. [FNA19, HC2]

Sp. Pl. 2: 914.

spotted knapweed

ssp. *australis* (A. Kern) Greuter [FNA, HC2]

Willdenowia 33(1): 56.

spotted knapweed

*Centaurea biebersteinii* DC., misapplied

*Centaurea maculosa* Lam. [HC], misapplied

*Centaurea paniculata* L., misapplied

*Centaurea stoebe* L. ssp. *micranthos* (S.G. Gmel. ex Gugler) Hayek [FNA19]

FNA19 includes a brief reference to this taxon in the text for *C. stoebe*. BC flora uses this interpretation, noting it is closely related to *C. paniculata*, but Chambers and Sundberg (2000) think the use of *C. bieb.* for our plants may be misapplied, and "needs further study".

*Centaurea trichocephala* M. Bieb. ex Willd. [FNA19, HC2]

Sp. Pl., ed. 4 [Willdenow] 3(3): 2286.

featherhead knapweed

FNA19: "A population of *Centaurea trichocephala* M. Bieberstein ex Willdenow (featherhead or hairy-head knapweed) was found in the late 1970s in a degraded pasture in eastern Washington (B. F. Roché and C. T. Roché 1991). A weed-control program was instituted, and the plants were successfully eradicated. Although it is apparently not established anywhere in North America, *C. trichocephala* is listed as a noxious weed in Oregon. These plants resemble *C. phrygia* in having elongate, pectinate-fringed phyllary appendages. In *C. trichocephala* the linear-filiform, featherlike appendages are much narrower than the phyllary bodies. Plants of the species spread by horizontal roots. According to Roché and Roché, *C. trichocephala* is apparently self-sterile; the Oregon plants spread clonally and formed no seeds."

*Centaurea ×varnensis* Velen. [HC2]

Fl. Bulg. 313.

hybrid diffuse knapweed, sand knapweed

(= *Centaurea diffusa* × *Centaurea stoebe* ssp. *micranthos*)

*Centaurea ×psammogena* G. Gayer

This name appears at the bottom of the description for *C. diffusa*. Both the International Plant Names Index (IPNI) and TROPICOS show this name published as "Centaurea psammogena Gayer". IPNI indicates that the name represents a taxon of hybrid origin.

*Centromadia* [FNA21, HC2]

Fl. Francisc. 4: 424. 1897.

spikeweed

*Centromadia pungens* (Hook. & Arn.) Greene [FNA21, HC2]

Man. Bot. San Francisco. 196.

common spikeweed

*Hemizonia pungens* (Hook. & Arn.) Torr. & A. Gray [HC]

ssp. *pungens* [FNA21, HC2]

Man. Bot. San Francisco. 196.

common spikeweed, western spikeweed

*Hemizonia pungens* (Hook. & Arn.) Torr. & A. Gray ssp. *septentrionalis* D.D. Keck

*Hemizonia pungens* (Hook. & Arn.) Torr. & A. Gray var. *pungens* [HC]

*Hemizonia pungens* (Hook. & Arn.) Torr. & A. Gray var. *septentrionalis* (D.D. Keck) Cronquist [HC]

Noxious in WA. FNA21: "Subspecies *pungens* is circumscribed broadly to include subsp. *maritima* and subsp. *septentrionalis* based on morphologic and molecular data (B. G. Baldwin, unpubl.). As treated here, *Centromadia pungens* subsp. *pungens* occurs widely in central and northern California, and it is putatively introduced in southwestern California and outside the state."

*Chaenactis* [FNA21, HC, HC2]

Prodri. 5: 659. 1836.

chaenactis, false-yarrow

*Chaenactis douglasii* (Hook.) Hook. & Arn. [FNA21, HC, HC2]

Bot. Beechey Voy. 354.

hoary chaenactis, hoary false-yarrow

var. *douglasii* [FNA21, HC, HC2]

Bot. Beechey Voy. 354.

dustymaidens, hoary false yarrow

*Chaenactis douglasii* (Hook.) H. & A var. *achilleaefolia* (H. & A.) A. Nels. [HC]

*Chaenactis douglasii* (Hook.) Hook. & Arn. var. *achilleifolia* (Hook. & Arn.) A. Gray

*Chaenactis douglasii* (Hook.) Hook. & Arn. var. *glandulosa* Cronquist [HC]

*Chaenactis douglasii* (Hook.) Hook. & Arn. var. *montana* M.E. Jones [HC]

*Chaenactis douglasii* (Hook.) Hook. & Arn. var. *rubricaulis* (Rydb.) Ferris

*Chaenactis pedicularia* Greene

*Chaenactis ramosa* Stockw. [HC]

FNA21: "Most of the diploid elements of var. *douglasii* are distinctive and are connected by a morphologically continuous series of polyploids (usually assigned to var. *achilleifolia*). Some diploid forms (including var. *rubricaulis* and *Chaenactis ramosa*) appear repeatedly and discontinuously in suitable habitats. In particular, forms named var. *montana* seem to arise wherever the species reaches sufficient elevation. Such populations have no historic or genetic cohesion to justify their recognition as a collective taxon, even though their reduced stature may become genetically fixed in each instance. (Variety *alpina*, recognized below with hesitation, may be just an extreme such case.)" Possibly not a syn, Kz99, BC & Chambers and Sundberg (2000) differ

*Chaenactis thompsonii* Cronquist [FNA21, HC, HC2]

Vasc. Pl. Pacif. N.W. 5: 123, fig. [p. 125].

Thompson's pincushion

FNA21: "Chaenactis thompsonii appears to be sister to C. evermannii; it is known from the mountains of central and northwestern Washington. The similar habits of C. thompsonii and C. ramosa (= C. douglasii var. douglasii) appear to result from convergent evolution in the distinctive habitat of their type localities (Wenatchee Mountains), not from a close genetic relationship as suggested by Cronquist."

***Chondrilla* [FNA19, HC, HC2]**

Sp. Pl. 2: 796. 1753; Gen. Pl. ed. 5, 348. 1754.  
gum-succory, skeletonweed

***Chondrilla juncea* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 796.  
hogbit, rush skeletonweed, gum succory

FNA19: "Chondrilla juncea is native to the Mediterranean region of Europe, North Africa, and Asia Minor. It is a weed in North America (not listed as noxious at the federal level). Its deep and extensive root system competes strongly for soil moisture and nutrients and makes control difficult because it helps the plants survive drought, cultivation, grazing, and most selective herbicides. The large, stiff branches and stems interfere with harvesting. The species is said to be "the most serious weed of Australian wheat-growing regions" (F. D. Panetta and J. Dodd 1987). It also infests millions of acres in California, Idaho, Oregon, and Washington. Chondrilla juncea is an obligate apomict; its seeds are formed by a parthenogenetic process (E. Battaglia 1949). Nevertheless, the species is highly variable in morphology and biochemical traits."

***Chrysothamnus* [FNA20, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 323. 1840.  
rabbit-brush  
(see also *Ericameria*)

***Chrysothamnus viscidiflorus* (Hook.) Nutt. [FNA20, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 324.  
green rabbit-brush

**ssp. *lanceolatus* (Nutt.) H.M. Hall & Clem. [FNA20, HC2]**

Publ. Carnegie Inst. Wash. 326: 181.  
sticky-leaf rabbitbrush, yellow rabbitbrush

*Chrysothamnus viscidiflorus* (Hook.) Nutt. var. *lanceolatus* (Nutt.) Greene [HC]

*Ericameria viscidiflora* (Hook.) L.C. Anderson ssp. *lanceolata* (Nutt.) L.C. Anderson

**ssp. *viscidiflorus* [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 324.  
sticky flowered rabbitbrush, sticky-leaf rabbitbrush, yellow rabbitbrush

*Chrysothamnus viscidiflorus* (Hook.) Nutt. ssp. *pumilus* (Nutt.) H.M. Hall & Clem.

*Chrysothamnus viscidiflorus* (Hook.) Nutt. ssp. *stenophyllus* (A. Gray) H.M. Hall & Clem.

*Chrysothamnus viscidiflorus* (Hook.) Nutt. var. *pumilus* (Nutt.) Jeps.

*Chrysothamnus viscidiflorus* (Hook.) Nutt. var. *stenophyllus* (A. Gray) H.M. Hall [HC]

*Chrysothamnus viscidiflorus* (Hook.) Nutt. var. *viscidiflorus* [HC]

*Ericameria viscidiflora* (Hook.) L.C. Anders. ssp. *viscidiflora*

*Ericameria viscidiflora* (Hook.) L.C. Anderson var. *stenophylla* (A. Gray) L.C. Anderson

***Cichorium* [FNA19, HC, HC2]**

Sp. Pl. 2: 813. 1753; Gen. Pl. ed. 5, 354. 1754.  
chicory

***Cichorium intybus* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 813.  
chicory, wild succory

***Cirsium* [FNA19, HC, HC2]**

Gard. Dict. Abr. ed. 4. vol. 1. 1754.  
thistle

*Cirsium arvense* (L.) Scop. [FNA19, HC, HC2]

Fl. Carniol. ed. 2. 2: 126.

Canadian thistle, creeping thistle

*Cirsium arvense* (L.) Scop. var. *arvense* [HC]

*Cirsium arvense* (L.) Scop. var. *horridum* Wimm. & Grab. [HC]

*Cirsium arvense* (L.) Scop. var. *mite* Wimm. & Grab.

***Cirsium brevifolium* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 421.

Palouse thistle

*Cirsium palousense* (Piper) Piper

***Cirsium brevistylum* Cronquist [FNA19, HC, HC2]**

Leafl. W. Bot. 7: 26.

clustered thistle, short-styled thistle

FNA19: "Cirsium brevistylum occurs in the coast ranges and adjacent coastal slope from southwestern British Columbia to southern California. In the Pacific Northwest its range extends inland to the northern Rocky Mountains of southern British Columbia, Idaho, and northwestern Montana, and the Blue and Walla Walla ranges of eastern Oregon. It is absent from the central and southern Cascade Range. In older literature the name *Cirsium edule* was widely misapplied to this species. A. Cronquist (1953) pointed out that the type of *C. edule* has corolla and style features quite different from those of the plants that had been called by that name and established the name *C. brevistylum*, based upon the notably short styles of this species. Hybrids of *C. brevistylum* with *C. edule* have been named *C. ×vancouverense* R. J. Moore & C. Frankton."

***Cirsium edule* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 420.

edible thistle

**var. *edule* [FNA19, HC2]**

edible thistle, Indian thistle, Macoun's thistle

*Carduus macounii* Greene

*Cirsium edule* Nutt. var. *macounii* (Greene) D.J. Keil [FNA19]

*Cirsium hallii* (A. Gray) M.E. Jones [HC]

*Cirsium macounii* (Greene) Petr.

**var. *wenatchense* D.J. Keil [FNA19, HC2]**

Sida. 21: 213.

Wenatchee thistle

FNA19: "Variety *wenatchense* is known only from the Wenatchee Mountains of central Washington. Little is known of its ecology." FNA key separates out this variety on the basis of the heads being mostly borne singly and peduncles 10?30 cm.

***Cirsium hookerianum* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 418.

Hooker's thistle, white thistle

***Cirsium inamoenum* (Greene) D.J. Keil [FNA19, HC2]**

Sida. 21: 214.

Greene's thistle

*Cirsium wallowense* M. Peck

**var. *inamoenum* [FNA19, HC2]**

Sida. 21: 214.

Greene's thistle

*Cirsium neomexicanum* A. Gray [FNA19, HC2], misapplied

*Cirsium subniveum* Rydb. [HC]

Known from a single collection in Garfield County. FNA19: "Plants of northeastern Oregon, southeastern Washington, and adjacent western Idaho often have large heads and densely tomentose

foliage. These were named *Cirsium wallowense* by Peck. Similar plants occur sporadically in other portions of the range of *Cirsium inamoenum* var. *inamoenum* and I chose not to recognize these northwestern populations as a third variety. Additional study might clarify the relationships of these plants."

***Cirsium remotifolium* (Hook.) DC. [FNA19, HC, HC2]**

Prod. 6: 655.

few-leaf thistle, Pacific fringed thistle, remote-leaved thistle, weak thistle

*Carduus remotifolius* Hook.

*Cirsium callilepis* (Greene) Jeps. [HC]

*Cirsium callilepis* (Greene) Jeps. var. *oregonense* (Petr.) J.T. Howell [HC]

*Cirsium remotifolium* (Hook.) DC. ssp. *remotifolium*

*Cirsium remotifolium* (Hook.) DC. var. *odontolepis* Petr. [FNA19]

*Cirsium remotifolium* (Hook.) DC. var. *remotifolium* [FNA19]

*Cirsium remotifolium* (Hook.) DC. var. *rivulare* Jeps. [FNA19]

FNA19: "Variety *remotifolium* occurs primarily west of the Cascade Range in Washington and Oregon and on coastal-facing slopes in northwestern California. Intermediates with var. *odontolepis* are known through much of that range." "Cirsium *remotifolium* occurs from the Coast Ranges and valleys of the Pacific Northwest to the western slopes of the Cascade and Klamath ranges, south in the California North Coast Ranges to the San Francisco Bay region. It is closely related to the *C. clavatum* complex of the Rocky Mountains region. Both have a similar growth habit and some forms variably express the character of broadly scarious, lacerate-toothed phyllary margins. Gray, in naming *Cnicus carlinoides* var. *americanus*, included as syntypes both California and Colorado specimens. F. Petrank (1917) treated both the West Coast plants and those of the Rocky Mountains as *Cirsium* subsect. *Americana*, recognizing *C. remotifolium* with several infraspecific taxa plus two other species, *C. callilepis* and *C. amblylepis* from the West Coast, and four additional species from the Rocky Mountains. A. Cronquist (1955) rejected Petrank's subspecies, treating *C. remotifolium* in a restricted sense, limited to plants of Washington and Oregon without dilated phyllary tips, and circumscribed *C. centaureae* broadly to include the Rocky Mountains and West Coast plants with dilated phyllary tips. Because of the frequent presence of dilated phyllary tips in *C. remotifolium* in the restricted sense, Cronquist acknowledged the likelihood of past introgression with *C. centaureae* in the broad sense. J. T. Howell (1960b) recognized three species: *Cirsium remotifolium*, *C. acanthodontum*, and *C. callilepis*, the latter with four varieties collectively corresponding to the West Coast representatives of *C. centaureae* (in the sense of Cronquist). Because of the great similarity of the various West Coast plants and their intergradation, I see no value in recognizing two or more species. The West Coast and Rocky Mountains plants are clearly related, but are separated by the Great Basin region and there is little chance of current genetic interchange. As is often the case with American *Cirsium*, genetic enrichment from past hybridization with other nearby species within their respective areas has likely been fertile ground for evolutionary diversification. Different species have contributed genes in the Pacific states and in the Rockies. I have chosen to recognize two geographically-based species complexes, each with intergrading races here treated as varieties. I treat the West Coast plants as *C. remotifolium* and the Rocky Mountains plants as *C. clavatum*."

***Cirsium scariosum* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 420.

elk thistle, meadow thistle

*Cirsium hookerianum* Nutt. var. *scariosum* (Nutt.) B. Boivin

*Cirsium magnificum* (A. Nelson) Petr. [HC]

*Cirsium scariosum* Nutt. var. *americanum* (A. Gray) D.J. Keil [FNA19]

*Cirsium scariosum* Nutt. var. *citrinum* (Petr.) D.J. Keil [FNA19]

*Cirsium scariosum* Nutt. var. *coloradense* (Rydb.) D.J. Keil [FNA19]

*Cirsium scariosum* Nutt. var. *congdonii* (R.J. Moore & Frankton) D.J. Keil [FNA19]

*Cirsium scariosum* Nutt. var. *robustum* D.J. Keil [FNA19]

*Cirsium scariosum* Nutt. var. *scariosum* [FNA19]

*Cirsium scariosum* Nutt. var. *thorneae* S.L. Welsh [FNA19]

*Cirsium scariosum* Nutt. var. *toiyabense* D.J. Keil [FNA19]

*Cirsium tiognum* (Congdon) Petr. var. *tiognum*

***Cirsium undulatum* (Nutt.) Spreng. [FNA19, HC, HC2]**

Syst. Veg. 3: 374.

wavy leaf thistle

*Cirsium undulatum* (Nutt.) Spreng. var. *undulatum*

FNA19: "Cirsium undulatum is widely distributed in the western half of North America from the dry plains and plateaus of the Pacific Northwest eastward across the Great Plains to Manitoba and the Dakotas and south to Texas, New Mexico, and northwestern Mexico. It occurs in scattered localities in the Rocky Mountains and northeastern Great Basin region. At least some of the few widely scattered records from the eastern United States are probably introductions. Cirsium undulatum is both widespread and variable. Plants of the Great Plains region tend to be low-growing with a few large heads and elongate corollas. Plants of the Pacific Northwest are usually taller and produce smaller, more numerous heads with shorter corollas. A detailed study of this species might reveal races worthy of recognition as infraspecific taxa. Wavyleaf thistle is listed by California as a noxious weed. However, most reports of Cirsium undulatum in California are based upon misidentifications of *C. canescens*. Cirsium undulatum is known to hybridize with *C. flodmanii*, *C. hookerianum*, and *C. scariosum* var. *coloradense*. J. T. Howell (1960b) reported that *C. undulatum* was suspected to hybridize with *C. brevifolium* in the Pacific Northwest."

*Cirsium vulgare* (Savi) Ten. [FNA19, HC, HC2]

Fl. Napol. 5: 209.

bull thistle, common thistle

*Carduus vulgaris* Savi

***Columbiadoria*** [FNA20, HC2]

Phytologia. 71: 249. 1991.

goldenweed

***Columbiadoria hallii*** (A. Gray) G.L. Nesom [FNA20, HC2]

Phytologia. 71: 249.

Columbia River daisy, Hall's goldenweed

*Haplopappus hallii* A. Gray [HC]

FNA20: "Columbiadoria hallii is known from the vicinity of the eastern Columbia River Gorge. It occurs also "at scattered stations [south] in the Cascades to the Calapooia Mountains," where the plants "are not precisely like the others, and may prove to be varietally distinct" (A. Cronquist 1955, p. 216)."

***Conyza*** [FNA20, HC, HC2]

Syn. Gen. Compos. 203. 1832.

conyza, horseweed

***Conyza bonariensis*** (L.) Cronquist [FNA20, HC, HC2]

Bull. Torrey Bot. Club. 70: 632.

South American conyza

(see also *Conyza sumatrensis* var. *sumatrensis*)

***Conyza canadensis*** (L.) Cronquist [FNA20, HC, HC2]

Bull. Torrey Bot. Club. 70: 632.

Canadian fleabane, horseweed

*Conyza canadensis* (L.) Cronquist var. *canadensis*

*Conyza canadensis* (L.) Cronquist var. *glabrata* (A. Gray) Cronquist

FNA20: "Conyza canadensis is thought to be native to North America and is now widely adventive, e.g., in South America, Europe, Asia, and Africa. Plants with stems glabrous and phyllaries red-tipped are sometimes treated as var. *pusilla*; similar plants with stems glabrous and phyllaries stramineous (not red-tipped) are sometimes treated as var. *glabrata*."

***Coreopsis*** [FNA21, HC, HC2]

Sp. Pl. 2: 907. 1753; Gen. Pl. ed. 5, 388. 1754.

coreopsis, tickseed

***Coreopsis grandiflora*** Hogg ex Sweet [FNA21, HC2]

Brit. Fl. Gard. 2: plate 175. 1826.

bigleaf tickseed

*Coreopsis grandiflora* Hogg ex Sweet × *Coreopsis lanceolata* L. [HC2]

*Coreopsis lanceolata* L. [FNA21, HC2]

Sp. Pl. 2: 908. 1753.

lance-leaved tickseed

Not in H&C. Recently (2008) collected in San Juan County.

*Coreopsis tinctoria* Nutt. [FNA21, HC, HC2]

J. Acad. Nat. Sci. Philadelphia. 2: 114.

calliopsis, Columbia coreopsis

*Coreopsis atkinsoniana* Douglas ex Lindl. [HC]

*Coreopsis tinctoria* Nutt. var. *atkinsoniana* (Douglas ex Lindl.) H.M. Parker ex E.B. Sm.

*Coreopsis tinctoria* Nutt. var. *tinctoria*

FNA21:"Coreopsis tinctoria is widely grown in public and residential gardens, and commercially (for cut flowers), and has become widely established in the flora area. As here circumscribed, Coreopsis tinctoria includes plants that others (without agreement among themselves) have treated as distinct species or infraspecific taxa: C. atkinsoniana (plants mostly 50?150+ cm, seldom branched from bases; cypselae 2.5?3 mm, "narrowly" winged; pappi 0.1?0.2 mm; mostly Idaho, Montana, Oregon, Washington), C. cardaminefolia (plants mostly 20?50 cm, seldom branched at bases; cypselae 2 mm, "narrowly to widely" winged; pappi 0 or 0.1?0.2 mm; mostly Arkansas, Kansas, Louisiana, Nebraska, Oklahoma, Texas), and C. tinctoria var. *similis* (plants mostly 10?30 cm, usually branched from bases; cypselae 2?3 mm, "widely" winged; pappi 0.2?1 mm; Texas and Mexico)."

*Cota* [FNA19, HC2]

Fl. Sicul. Syn. 2: 866. 1845.

chamomile

*Cota austriaca* (Jacq.) Sch. Bip. [HC2]

Oesterr. Bot. Wochensbl. 4: 155.

Austrian chamomile

*Anthemis austriaca* Jacq. [Stace 1997]

*Cota tinctoria* (L.) J. Gay ex Guss. [FNA19, HC2]

Fl. Sicul. Syn. 2: 867.

golden chamomile, yellow chamomile

*Anthemis tinctoria* L. [HC, Stace 1997]

This species is not included in FNA19.

*Cotula* [FNA19, HC, HC2]

Sp. Pl. 2: 891. 1753; Gen. Pl. ed. 5, 380. 1754.

cotula

*Cotula coronopifolia* L. [FNA19, HC, HC2]

Sp. Pl. 2: 892.

brass buttons, common brass buttons

*Crepis* [FNA19, HC, HC2]

Sp. Pl. 2: 805. 1753; Gen. Pl. ed. 5, 350. 1754.

hawksbeard

(see also *Askellia*)

*Crepis acuminata* Nutt. [FNA19, HC, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 437.

long-leaved hawksbeard, tapertip hawksbeard

*Crepis acuminata* Nutt. ssp. *acuminata* [HC]

*Crepis atribarba* A. Heller [FNA19, HC2]

Bull. Torrey Bot. Club. 26: 314.

slender hawksbeard

*Crepis atrabarba* Heller [HC], orthographic variant  
*Crepis atrabarba* A. Heller ssp. *atrabarba* [HC], orthographic variant  
*Crepis atrabarba* A. Heller ssp. *originalis* Babc. & Stebbins [HC], orthographic variant  
*Crepis atrabarba* A. Heller ssp. *atribarba*  
*Crepis atrabarba* A. Heller ssp. *originalis* (Babc. & Stebbins) Babc. & Stebbins

FNA19: "Crepis atrabarba is generally recognized by the deeply pinnately lobed leaves with linear lobes, fine tomentulose indument on stems and leaves, setose phyllaries, and dark green, strongly ribbed cypselae. It is a variable mixture that includes polyploid, apomictic forms and hybrids with *C. acuminata* and other species. The typical form is recognized by its short stature, narrow pinnately lobed, tomentulose leaves, stems with 3?10 heads, and phyllaries with scattered, black, eglandular setae. Larger, more robust forms with stems 30?70 cm, 10?30+ heads, narrower involucres, and few or no black setae have been recognized as subsp. *originalis*. The latter was considered by E. B. Babcock (1947) to represent the original diploid form of the species; it is difficult to distinguish in practice."

***Crepis bakeri* Greene [FNA19, HC, HC2]**

Erythea. 3: 73.  
Baker's hawksbeard

*Crepis bakeri* Greene ssp. *bakeri* [FNA19, HC]  
*Crepis bakeri* Greene ssp. *cusickii* (Eastw.) Babc. & Stebbins [FNA19]  
*Crepis bakeri* Greene ssp. *idahoensis* Babc. & Stebbins [FNA19, HC]

FNA19: "Crepis bakeri is generally recognized by the low stature, dense rosettes of pinnately lobed leaves with coarsely dentate lobes, tomentose stems and leaves, stipitate-glandular hairs distally on stems, relatively large involucres, and densely flowered heads. It is considered closely related to *C. occidentalis*. Three somewhat weakly defined subspecies were recognized by E. B. Babcock (1947)."

***Crepis barbigera* Leiberg ex Coville [FNA19, HC, HC2]**

Contr. U.S. Natl. Herb. 3: 565, plate 26.  
bearded hawksbeard

FNA19: "Crepis barbigera is recognized by its relatively tall stature, deeply pinnately lobed leaves, tomentulose stems, and phyllaries with coarse, green, eglandular setae. It is a complex of polyploid, apomictic forms, combining characteristics of *C. atrabarba*, *C. acuminata*, and *C. modocensis*, from which the species is presumed to have been derived by intercrossing (E. B. Babcock 1947)."

***Crepis capillaris* (L.) Wallr. [FNA19, HC, HC2]**

Linnaea. 14: 657.  
smooth hawksbeard

*Crepis capillaris* (L.) Wallr. var. *capillaris*

FNA19: "Crepis capillaris is recognized by its shallow root system, dense rosettes of coarsely dentate or pinnately lobed leaves, erect slender stems, auriculate-based caudine leaves, relatively small heads, phyllaries with double rows of black setae, and fluffy white pappi. It is weedy and can become a serious lawn pest. It is one of only three species of Crepis with  $2n = 6$ ; E. B. Babcock (1947) considered it to be advanced in the genus." Reports of var. *agrestis* Atkinson & Sharpe (1993) are not supported by a specimen.

***Crepis intermedia* A. Gray [FNA19, HC, HC2]**

Syn. Fl. N. Amer. 1(2): 432.  
gray hawksbeard, intermediate hawksbeard, limestone hawksbeard

***Crepis modocensis* Greene [FNA19, HC, HC2]**

Erythea. 3: 48.  
low hawksbeard, Modoc hawksbeard

*Crepis modocensis* Greene ssp. *glareosa* (Piper) Babc. & Stebbins [FNA19]  
*Crepis modocensis* Greene ssp. *modocensis* [FNA19, HC]  
*Crepis modocensis* Greene ssp. *rostrata* (Coville) Babc. & Stebbins [FNA19, HC]  
*Crepis modocensis* Greene ssp. *subacaulis* (Kellogg) Babc. & Stebbins [FNA19]  
*Crepis rostrata* Coville

***Crepis nicaeensis* Balbis ex Pers. [FNA19, HC, HC2]**

Syn. Pl. 2: 376.

French hawksbeard, Turkish hawksbeard

FNA19:"*Crepis nicaeensis* is distinguished by the annual or biennial habit, shallow root system, hispid stems, and glabrate phyllaries enclosing outer cypselae. It is similar in habit to *C. biennis*, which differs in its larger heads and 13?20-ribbed cypselae; it is considered closely related to *C. capillaris* (E. B. Babcock 1947)."

***Crepis occidentalis* Nutt. [FNA19, HC, HC2, JPM]**

J. Acad. Nat. Sci. Philadelphia. 7: 29.  
western hawksbeard

*Crepis occidentalis* Nutt. ssp. *conjugata* Babc. & Stebbins [FNA19, HC]  
*Crepis occidentalis* Nutt. ssp. *costata* (A. Gray) Babc. & Stebbins [FNA19, HC]  
*Crepis occidentalis* Nutt. ssp. *occidentalis* [FNA19, HC]  
*Crepis occidentalis* Nutt. ssp. *pumila* (Rydb.) Babc. & Stebbins [FNA19, HC]  
*Crepis occidentalis* Nutt. var. *costata* A. Gray

Taxonomy follows Stebbins in Jepson Manual in not recognizing subspecies in species of *Crepis* that are largely apomictic. FNA19 does recognize four subspecies. FNA19:"*Crepis occidentalis* is recognized by the old, brown leaf bases persisting on caudices, by stems, leaves, and phyllaries gray-tomentose, and by loose, corymbiform arrays with relatively few, relatively large heads. It is widespread and polymorphic. Some specimens have coarse setae or black, stipitate glands on the phyllaries in addition to the tomentose indument, the stipitate glands sometimes extending proximally on stems. Four intergrading subspecies were recognized by E. B. Babcock (1947). The sexual diploid forms are found in subsp. *occidentalis* and occur in northern California and adjacent Nevada. The other subspecies are polyploid and apomictic (Babcock)."

***Crepis runcinata* (E. James) Torr. & A. Gray [FNA19, HC, HC2]**

Fl. N. Amer. 2: 487.  
dandelion hawksbeard, meadow hawksbeard

**ssp. *runcinata* [FNA19, HC, HC2]**

Fl. N. Amer. 2: 487.  
dandelion hawksbeard, meadow hawksbeard

*Crepis runcinata* (E. James) Torr. & A. Gray ssp. *glauca* (Nutt.) Babc. & Stebbins [FNA19, HC]  
*Crepis runcinata* (E. James) Torr. & A. Gray ssp. *hispidulosa* (Howell ex Rydb.) Babc. & Stebbins [FNA19, HC]  
*Crepis runcinata* (E. James) Torr. & A. Gray ssp. *imbricata* Babc. & Stebbins [FNA19]  
*Crepis runcinata* (E. James) Torr. & A. Gray var. *hispidulosa* Howell ex Rydb.

***Crepis setosa* Haller f. [FNA19, HC, HC2]**

Arch. Bot. (Leipzig). 1(2): 1.  
bristly hawksbeard, rough hawksbeard

FNA19:"*Crepis setosa* is recognized by its annual habit, shallow roots, coarsely setose stems, leaves, and involucres, the relatively large runcinate leaves, sagittate-laciniate cauline leaves, finely beaked cypselae, and white, fine pappus bristles."

***Crepis tectorum* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 807.  
annual hawksbeard, narrow leaf hawksbeard, rooftop hawksbeard

***Crocidium* [FNA20, HC, HC2]**

Fl. Bor.-Amer. 1: 335, plate 118. 1834.  
crocidium, spring-gold

***Crocidium multicaule* Hook. [FNA20, HC, HC2]**

Fl. Bor.-Amer. 1: 335, plate 118.  
gold-star, spring-gold

***Crupina* [FNA19, HC, HC2]**

Ann. Mus. Natl. Hist. Nat. 16: 157. 1810.  
crupina

*Crupina vulgaris* Pers. ex Cass. [FNA19, HC, HC2]

Dict. Sci. Nat. ed 2. 12: 68.

bearded creeper, crupina

***Cyclachaena* [FNA21, HC2]**

Index Seminum (Frankfurt). 1836: 4. 1838.

marsh-elder

*Cyclachaena xanthiiifolia* (Nutt.) Fresenius [FNA21, HC2]

Index Seminum (Frankfurt). 1836: 4.

carelessweed, burweed marsh-elder, tall marsh-elder

*Iva xanthifolia* Nutt. [HC]

*Iva xanthiiifolia* Nutt.

FNA21: "Cyclachaena xanthiiifolia is thought to be native to North American prairies and is evidently adventive east of the Mississippi River and in western states. It was recorded once from California as a weed in commercially grown carrots (specimen in CAS)."

***Dieteria* [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 300. 1840.

tansyaster

*Dieteria canescens* (Pursh) Nutt. [FNA20, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 300.

hoary-aster

*Machaeranthera canescens* (Pursh) A. Gray [HC]

var. *incana* (Lindl.) D.R. Morgan & R.L. Hartm. [FNA20, HC2]

Sida. 20: 1396. 2003.

hoary-aster, tall hoary-aster

*Aster attenuatus* (Howell) Frye & Rigg ex M. Peck, illegitimate name

*Dieteria incana* (Lindl.) Torr. & A. Gray

*Diplopappus incanus* Lindl.

*Machaeranthera canescens* (Pursh) A. Gray var. *incana* (Lindl.) A. Gray

*Machaeranthera incana* (Lindl.) Greene

***Doronicum* [FNA20, HC2]**

Sp. Pl. 2: 885. 1753; Gen. Pl. ed. 5, 377. 1754.

leopard's bane

*Doronicum willdenowii* (Rouy) A.W.Hill [New Flora of the British Isles, 2nd Edition]

Index Kew. Suppl. 6: 71.

Willdenow's leopard-bane

(= *Doronicum pardalianches* x *Doronicum plantagineum*)

Collected in 2022 in King County.

***Eatonella* [FNA21, HC, HC2]**

Proc. Amer. Acad. Arts. 19: 19. 1883.

Eatonella

*Eatonella nivea* (D.C. Eaton) A. Gray [FNA21, HC, HC2]

Proc. Amer. Acad. Arts. 19: 19.

white Eatonella, white false tickhead

*Burielia nivea* D.C. Eaton

FNA21 does not show this species occurring in WA, however there are specimens from WA at WTU.

***Echinops* [FNA19, HC, HC2]**

Sp. Pl. 2: 814. 1753; Gen. Pl. ed. 5, 356. 1754.

globe-thistle

ssp. *ruthenicus* (M. Bieb.) Nyman [FNA19, HC2]

Consp. Fl. Eur. 2: 399. 1879.

southern globe thistle

*Echinops ruthenicus* M. Bieb. [HC]

No specimens of this taxon are known from Washington, despite being reported in WA by Abrams and Kartesz. Until a specimen is produced from WA, this taxon will not be considered part of the flora.

***Erechtites* [FNA20, HC, HC2]**

Fl. Ludov. 65. 1817.

burnweed, fireweed

*Erechtites minus* (Poir.) DC. [FNA20, HC2]

Prodri. 6: 437.

Australian burnweed, toothed coast burnweed

*Erechtites minima* (Poir.) DC. [HC], orthographic variant

*Erechtites prenanthoides* (A. Richardson) DC.

*Senecio minus* Poir. [JPM2]

***Ericameria* [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 318. 1840.

goldenbrush, rabbit-brush

*Ericameria bloomeri* (A. Gray) J.F. Macbr. [FNA20, HC2]

Contr. Gray Herb. 56: 36.

Bloomer's goldenweed, rabbitbrush goldenweed

*Haplopappus bloomeri* A. Gray [HC]

*Haplopappus bloomeri* A. Gray var. *angustatus* A. Gray

*Haplopappus bloomeri* A. Gray var. *bloomeri*

*Haplopappus bloomeri* A. Gray var. *sonnei* Greene

Intergrades with *E. greenei*.

*Ericameria greenei* (A. Gray) G.L. Nesom [FNA20, HC2]

Phytologia. 68: 153.

Greene's goldenweed, Greene's heath goldenweed

*Haplopappus bloomeri* A. Gray var. *greenei* (A. Gray) Cronquist

*Haplopappus greenei* A. Gray [HC]

*Haplopappus greenei* A. Gray var. *greenei*

*Haplopappus greenei* A. Gray var. *mollis* A. Gray

*Ericameria nauseosa* (Pall. ex Pursh) G.L. Nesom & G.I. Baird [FNA20, HC2]

Phytologia. 75: 84. 1993.

common rabbit-brush

*Chrysanthemum nauseosum* (Pall. ex Pursh) Britton [HC]

var. *nana* (Cronquist) G.L. Nesom & G.I. Baird [FNA20, HC2]

Phytologia. 75: 87.

little rabbitbrush

*Chrysanthemum nauseosum* (Pall. ex Pursh) Britton ssp. *nanus* (Cronquist) D.D. Keck

*Chrysanthemum nauseosum* (Pall. ex Pursh) Britton var. *nanus* Cronquist [HC]

var. *speciosa* (Nutt.) G.L. Nesom & G.I. Baird [FNA20, HC2]

Phytologia. 75: 87.

rubber rabbitbrush

*Chrysanthemum nauseosum* (Pall. ex Pursh) Britton ssp. *albicaulis* (Nutt.) H.M. Hall & Clem.

*Chrysanthemum nauseosum* (Pall. ex Pursh) Britton ssp. *speciosus* (Nutt.) H.M. Hall & Clem.

*Chrysanthemum nauseosum* (Pall. ex Pursh) Britton var. *albicaulis* (Nutt.) Rydb. [HC]

*Chrysanthemum nauseosum* (Pall. ex Pursh) Britton var. *speciosus* (Nutt.) H.M. Hall

***Ericameria resinosa*** Nutt. [FNA20, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 319.

Columbia goldenweed, Columbian heath goldenweed

*Haplopappus resinosus* (Nutt.) A. Gray [HC]

***Erigeron*** [FNA20, HC, HC2]

Sp. Pl. 2: 863. 1753; Gen. Pl. ed. 5, 371. 1754.

daisy, erigeron, fleabane

***Erigeron acris*** L. [FNA20, HC, HC2]

Sp. Pl. 2: 863.

bitter fleabane

(see also *Erigeron elatus*, *Erigeron nivalis*)

**var. *kamtschaticus* (DC.) Herder** [FNA20, HC2]

Bull. Soc. Imp. Naturalistes Moscou. 38: 392.

Kamchatka bitter fleabane

*Erigeron acris* L. ssp. *politus* (Fr.) Schinz & R. Keller

*Erigeron acris* L. var. *asteroides* (Andrz. ex Besser) DC. [HC]

*Trimorpha acris* (L.) Gray var. *asteroides* (Andrz. ex Besser) G.L. Nesom

***Erigeron aliceae*** Howell [FNA20, HC, HC2]

Fl. N.W. Amer. 317.

Alice's fleabane, Eastwood's fleabane

***Erigeron annuus*** (L.) Pers. [FNA20, HC, HC2]

Syn. Pl. 2: 431.

annual fleabane, eastern daisy fleabane, sweet scabrous fleabane

*Aster annuus* L.

*Erigeron annuus* (L.) Pers. var. *discoideus* (Vicq. & J. Rouss.) Cronquist

***Erigeron aureus*** Greene [FNA20, HC, HC2]

Pittonia. 2: 16.

golden fleabane

*Erigeron aureus* Greene var. *acutifolius* Raup

FNA20: "Erigeron aureus var. *acutifolius* has leaves apically acute (versus rounded to broadly obtuse, sometimes emarginate, in the typical form) and is known only from the type locality, a peat bog in British Columbia (Peace River District). It was not listed or otherwise recognized in a recent flora of that province (G. W. Douglas et al. 1998?2002, vol. 1). *Erigeron xanthurii* B. Boivin was described as "sp. nov." and was noted to have originated as a hybrid between *E. acris* and *E. aureus*. It was treated by E. H. Moss and J. G. Packer (1983) as a hybrid. Specimens cited by Boivin are from widely separated localities in southwestern British Columbia and adjacent Alberta. It was included at specific rank in the treatment by A. C. Budd et al. (1987) but not by H. J. Scoggan (1978?1979, part 4) or G. W. Douglas et al. (1998?2002, vol. 1)."'

***Erigeron basalticus*** Hoover [FNA20, HC, HC2]

Leafl. W. Bot. 4: 40.

basalt fleabane

***Erigeron bloomeri*** A. Gray [FNA20, HC, HC2]

Proc. Amer. Acad. Arts. 6: 540.

Bloomer's fleabane, scabland fleabane

**var. *bloomeri*** [FNA20, HC, HC2]

Proc. Amer. Acad. Arts. 6: 540.

Bloomer's fleabane, scabland fleabane

***Erigeron caespitosus*** Nutt. [FNA20, HC, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 307.

tufted fleabane

FNA20: "Erigeron caespitosus as recognized here is highly variable and perhaps justifiably could be

divided into more than one taxon. Plants at lower elevations tend to produce tall stems branching above the middle and long, white rays. At higher elevations, especially in Idaho, western Montana, Utah, and Wyoming, stems tend to be shorter and simple and the rays commonly are blue to violet. In the Bitterroot Mountains (Ravalli and Deerlodge counties, Montana), short-stemmed, blue-rayed plants also have strigose caudine vestiture (in contrast to typically deflexed-hirtellous stems); these vestiture variants occur in the same area with plants apparently similar in all other features. Strigose populational variants also occur in Saskatchewan and Yukon, and *E. abajoensis*, largely distinguished by strigose caudine vestiture, might be considered a regional variant of *E. caespitosus*. In eastern Idaho and southwestern Montana, plants of *E. caespitosus* are commonly encountered with caudine leaves obovate and distinctly subclasping. Plants with strongly 3-nerved basal leaves occur in Carbon and Gallatin counties, Montana.

***Erigeron chrysopsisid*** A. Gray [FNA20, HC, HC2]

Syn. Fl. N. Amer. 1(2): 210.

dwarf yellow fleabane

**var. *chrysopsisid*** [FNA20, HC, HC2]

In A. Gray et al., Syn. Fl. N. Amer. 1(2): 210.

dwarf yellow fleabane, golden fleabane

*Erigeron chrysopsisid* A. Gray ssp. *chrysopsisid* [HC]

***Erigeron compositus*** Pursh [FNA20, HC, HC2]

Fl. Amer. Sept. 2: 535. 1813.

cutleaf daisy, dwarf mountain fleabane, fernleaf fleabane, trifid mountain fleabane

*Erigeron compositus* Pursh var. *compositus* [HC]

*Erigeron compositus* Pursh var. *discoideus* A. Gray [HC]

*Erigeron compositus* Pursh var. *glabratus* Macoun [HC]

*Erigeron trifidus* Hook. [HC2], misapplied

FNA20: "Correlations among ploidal level, breeding systems, and morphologic variation have been studied in detail in *Erigeron compositus*. Five informally designated population systems of diploids are geographically restricted (all of the northwestern United States and adjacent Canada) and primarily sexual, compared to the polyploids, which are agamospermous and apparently of hybrid origin, at least in some cases (R. D. Noyes et al. 1995; Noyes and D. E. Soltis 1996). Reduction in ray floret laminae usually is correlated with polyploidy. Plants with 1-ternately lobed leaves have been identified as var. *glabratus*, an element of variation that does not have a geographic pattern. Among closely related species, *Erigeron compositus* is the only one that produces strongly thickened caudex branches; occasional collections show a tendency toward the slender, loose branches characteristic of the other species."

***Erigeron corymbosus*** Nutt. [FNA20, HC, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 308.

foothill fleabane, longleaf fleabane

***Erigeron davisii*** (Cronquist) G.L. Nesom [FNA20, HC2]

Sida. 21: 22.

Davis's daisy, Davis's fleabane

*Erigeron engelmannii* A. Nelson var. *davisii* (Cronquist) Cronquist [HC]

Specimens from southeastern WA.

***Erigeron disparipilus*** Cronquist [FNA20, HC, HC2]

Brittonia. 6: 194.

Snake River fleabane, white cushion fleabane

***Erigeron divergens*** Torr. & A. Gray [FNA20, HC, HC2]

Fl. N. Amer. 2: 175.

*Erigeron divergens* Torr. & A. Gray var. *divergens* [HC2]

FNA20: "Polyploidy and agamospermy apparently are common in *Erigeron divergens* and contribute to the variability and, probably to some extent, the polymorphism characteristic of this species. Diploids appear to be scattered through the range of the species, at least in its southern part."

***Erigeron eatonii*** A. Gray [FNA20, HC, HC2]

Notes Compositae. 91. 1880.

Eaton's fleabane

var. *vilosus* (Cronquist) Cronquist [FNA20, HC, HC2]

Vasc. Pl. Pacif. N.W. 5: 175.

Eaton's shaggy fleabane

*Erigeron eatonii* A. Gray ssp. *vilosus* Cronquist

***Erigeron elatus* (Hook.) Greene [FNA20, HC2]**

Pittonia. 3: 164.

swamp fleabane

*Erigeron acris* L. var. *elatus* (Hook.) Cronquist [HC]

*Trimorpha acris* (L.) A. Gray var. *elatus* (Hook.) G.L. Nesom

*Trimorpha elata* (Hook.) G.L. Nesom

Reported from WA in FNA. Currently no specimens exist in Pacific Northwest herbaria.

***Erigeron filifolius* (Hook.) Nutt. [FNA20, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 308.

Peck's threadleaf fleabane

*Erigeron filifolius* (Hook.) Nutt. var. *filifolius* [HC]

*Erigeron filifolius* (Hook.) Nutt. var. *robustior* M. Peck [HC]

FNA20: "The densely white-strigose stem bases, linear-filiform leaves relatively unreduced distally, and relatively few heads with coiling, usually blue rays are distinctive for *Erigeron filifolius*. Proximal leaves are not clustered as a basal rosette; they are inserted on closely spaced nodes that are slightly more separated distally. Plants identified as var. *robustior* (with more ray florets, fewer heads, and thicker stems, centered in Oregon and Washington) intergrade with the typical form and apparently are separated arbitrarily."

***Erigeron flettii* G.N. Jones [FNA20, HC, HC2]**

Bot. Surv. Olympic Penins. 244.

Flett's fleabane, Olympic Mt. fleabane

FNA20: "*Erigeron flettii* differs from *E. grandiflorus* in having fewer, wider, consistently white rays, broadly spatulate basal leaves with bases constricted into narrow petioles longer than the blades and apices rounded or obtuse, less dense involucral vestiture, and strongly barbellate pappus bristles. It is known only from the Olympic Mountains."

***Erigeron glacialis* (Nutt.) A. Nelson [FNA20, HC2]**

Bot. Gaz. 37: 270.

glacier fleabane

var. *glacialis* [FNA20, HC2]

Bot. Gaz. 37: 270.

subalpine fleabane

*Erigeron peregrinus* (Banks ex Pursh) Greene ssp. *callianthemus* (Greene) Cronquist [HC]

*Erigeron peregrinus* (Banks ex Pursh) Greene ssp. *callianthemus* (Torr. & A. Gray) Cronquist

*Erigeron peregrinus* (Banks ex Pursh) Greene var. *angustifolius* (A. Gray) Cronquist [HC]

*Erigeron peregrinus* (Banks ex Pursh) Greene var. *callianthemus* (Greene) Cronquist

*Erigeron peregrinus* (Banks ex Pursh) Greene var. *eucallianthemus* Cronquist [HC]

*Erigeron peregrinus* (Banks ex Pursh) Greene var. *scaposus* (Torr. & A. Gray) Cronquist [HC]

Plants previously included within *E. peregrinus* (except *E. peregrinus* var. *thompsonii*) are now included within *E. glacialis* var. *glacialis*.

***Erigeron howellii* (A. Gray) A. Gray [FNA20, HC, HC2]**

Syn. Fl. N. Amer. 1(2): 209.

Howell's fleabane

*Erigeron salsuginosus* (Richardson ex R. Br.) A. Gray var. *howellii* A. Gray

***Erigeron inornatus* (A. Gray) A. Gray [FNA20, HC, HC2]**

Notes Compositae. 88.

California rayless fleabane

var. *inornatus* [FNA20, HC, HC2]

Notes Compositae. 88.

California rayless fleabane

***Erigeron leibergii* Piper [FNA20, HC, HC2]**

Bull. Torrey Bot. Club. 28: 41.

Leiberg's fleabane

***Erigeron linearis* (Hook.) Piper [FNA20, HC, HC2]**

Contr. U.S. Natl. Herb. 11: 567.

desert yellow daisy, lineleaf fleabane

*Diplopappus linearis* Hook.

*Erigeron peucephyllus* A. Gray

***Erigeron lonchophyllus* Hook. [FNA20, HC, HC2]**

Fl. Bor.-Amer. 2: 18.

short-rayed fleabane, spear-leaved fleabane

*Erigeron acris* L. ssp. *racemosus* (Nutt.) Clem. & E.G. Clem.

*Trimorpha lonchophylla* (Hook.) G.L. Nesom

***Erigeron nivalis* Nutt. [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 311.

northern daisy

*Erigeron acris* L. ssp. *debilis* (A. Gray) Piper

*Erigeron acris* L. var. *debilis* A. Gray [HC]

*Trimorpha acris* (L.) Gray var. *debilis* (A. Gray) G.L. Nesom

FNA20: "Erigeron nivalis usually has been treated as an infra-specific taxon within *E. acris*; the two are broadly sympatric in the northwestern United States and Canada without obvious intergrades. Both occur over a wide range of elevations and in similar habitats. *Erigeron nivalis* probably occurs in Nevada; it has not been taxonomically distinguished there. *Erigeron scotteri* was regarded by E. H. Moss and J. G. Packer (1983) as a synonym of *E. acris* (presumably var. *debilis* = *E. nivalis*; the heads are relatively small and borne singly)."

***Erigeron oreganus* A. Gray [FNA20, HC, HC2]**

Proc. Amer. Acad. Arts. 19: 2.

Gorge fleabane, Oregon fleabane

***Erigeron peregrinus* (Banks ex Pursh) Greene [FNA20, HC, HC2]**

Pittonia. 3: 166.

wandering fleabane

(see also *Erigeron glacialis*)

var. *peregrinus* [FNA20, HC2]

Pittonia. 3: 166.

wandering daisy

*Erigeron peregrinus* (Banks ex Pursh) Greene ssp. *peregrinus* [HC]

*Erigeron peregrinus* (Banks ex Pursh) Greene var. *dawsonii* Greene [HC]

*Erigeron peregrinus* (Banks ex Pursh) Greene var. *peregrinus* [FNA20, HC2], misapplied

Recently (2019) collected in northern Washington.

var. *thompsonii* (S.F. Blake ex J.W. Thomp.) Cronquist [FNA20, HC, HC2]

Brittonia. 6: 144.

Thompson's wandering fleabane

*Erigeron thompsonii* S.F. Blake ex J.W. Thomp.

***Erigeron philadelphicus* L. [FNA20, HC, HC2]**

Sp. Pl. 2: 863.

Philadelphia fleabane

var. *glaber* J.K. Henry [FNA20, HC2]

Ottawa Naturalist. 31: 57.

smooth Philadelphia fleabane

**var. *philadelphicus* [FNA20, HC2]**

Sp. Pl. 2: 863.

Philadelphia fleabane

***Erigeron piperianus* Cronquist [FNA20, HC, HC2]**

Brittonia. 6: 197.

Piper's fleabane

***Erigeron poliospermus* A. Gray [FNA20, HC, HC2]**

Syn. Fl. N. Amer. 1(2): 210.

hairy-seeded daisy, cushion fleabane

**var. *cereus* Cronquist [FNA20, HC, HC2]**

Brittonia. 6: 194.

Kittitas fleabane

FNA20: "Variety cereus grows in Chelan, Douglas, Grant, and Kittitas counties, apparently occurring as an enclave within the range of the typical variety. The vestiture and elongate proximal internodes of var. cereus are distinctive even within the group of species most closely related to *Erigeron poliospermus*; intergrades with typical *E. poliospermus* in vestiture and habit appear to be relatively common. Analogous variants occur within *E. concinnus* and are recognized at varietal rank."

**var. *poliospermus* [FNA20, HC, HC2]**

Brittonia. 6: 194.

hairy seeded daisy, cushion fleabane

***Erigeron pumilus* Nutt. [FNA20, HC, HC2]**

Gen. N. Amer. Pl. 2: 147.

shaggy fleabane

**var. *intermedius* (Cronquist) S.L. Welsh [FNA20, HC2]**

Brittonia. 6: 180.

shaggy fleabane

*Erigeron pumilus* Nutt. ssp. *intermedius* Cronquist [HC]

*Erigeron pumilus* Nutt. var. *euintermedius* Cronquist [HC]

*Erigeron pumilus* Nutt. var. *gracilior* Cronquist [HC]

FNA20: "Variety *gracilior* was described by A. Cronquist (1947) as "plants slender, the larger stems either not more than 1.5 mm thick near the base or bearing fewer than 5 heads." Such plants occur mostly in the southern part of the variety's range (mostly Idaho and Oregon, some in Washington); while the gracile tendency seems real, many arbitrary identifications must be made if two taxa are recognized."

**var. *pumilus* [FNA20, HC2]**

Gen. N. Amer. Pl. 2: 147.

shaggy fleabane

*Erigeron pumilus* Nutt. ssp. *pumilus* [HC]

***Erigeron salishii* G.W. Douglas & Packer [FNA20, HC2]**

Canad. J. Bot. 66: 414, fig. 1.

Salish fleabane, star peak fleabane

Often confused with *E. trifidus*, or *E. compositus*.

***Erigeron speciosus* (Lindl.) DC. [FNA20, HC, HC2]**

in A. P. de Candolle and A. L. P. P. de Candolle, Prodr. 5: 284

showy daisy, showy fleabane, splendid fleabane

*Erigeron speciosus* (Lindl.) DC. var. *macranthus* (Nutt.) Cronquist [HC]

*Erigeron speciosus* (Lindl.) DC. var. *speciosus* [HC]

*Erigeron subtrinervis* Rydb. ex Porter & Britton ssp. *conspicuus* (Rydb.) Cronquist

*Erigeron subtrinervis* Rydb. ex Porter & Britton var. *conspicuus* (Rydb.) Cronquist [HC]

*Stenactis speciosa* Lindl.

Taxonomy follows Flora of North America not recognizing varieties; FNA 20 (Nesom): "Plants glabrous and glandular on the phyllaries, stems, and leaves have been recognized as var. <i>macranthus</i>; they intergrade with hairier forms and do not show a coherent geographic pattern."

***Erigeron strigosus* Muhl. ex Willd. [FNA20, HC, HC2]**

Sp. Pl., ed. 4 [Willdenow] 3(3): 1956.

branched fleabane, daisy fleabane

**var. *septentrionalis* (Fernald & Wiegand) Fernald [FNA20, HC, HC2]**

Rhodora. 44: 340.

prairie fleabane

*Erigeron strigosus* Muhl. ex Willd. ssp. *septentrionalis* (Fernald & Wiegand) Wagenitz

FNA considers this taxon to be native. FNA20: "The distributional data given here are highly provisional; the author has not attempted to sort this taxon accurately, if it can be. According to A. Cronquist (1947), var. *septentrionalis* is scattered mostly in the northern half of the continent and (1994) is "found chiefly in New England and adjacent Canada." It is "morphologically transitional" to *Erigeron annuus* (A. Cronquist 1994) and, as implied by the synonymy, may be more appropriately treated as part of *E. annuus* (D. Frey et al. 2003). As noted by Fernald in the original description, the stems may be nearly glabrous or lightly hispid, in contrast to the more densely strigose to strigillose ones of typical *E. strigosus*." Cronquist (1955, Vol 5) believes western plants called this are hybrids (*E. annuus* x *strigosus*), and var. *septentrionalis* is native in e US.

**var. *strigosus* [FNA20, HC, HC2]**

Sp. Pl. 3: 1956.

branching fleabane, daisy fleabane

*Erigeron annuus* (L.) Pers. ssp. *strigosus* (Muhl. ex Willd.) Wagenitz

FNA considers this taxon native. Introduced (in OR, Chambers and Sundberg (2000), uncertain if native in BC (Douglas et al. 1998 Reported in WA by Creso (1984)

***Erigeron subtrinervis* Rydb. ex Porter & Britton [FNA20, HC, HC2]**

Mem. Torrey Bot. Club. 5: 328.

three-veined fleabane

FNA20: "Erigeron subtrinervis is variable in vestiture, perhaps reflecting gene exchange with *E. speciosus*. *Erigeron speciosus* var. *mollis* (A. Gray) S. L. Welsh may be a recurrent hybrid; it is identified here within *E. subtrinervis*."

***Eriophyllum* [FNA21, HC, HC2]**

Gen. Sp. Pl. 28. 1816.

woolly sunflower

***Eriophyllum lanatum* (Pursh) J. Forbes [FNA21, HC, HC2]**

Hort. Woburn. 183. 1833.

Oregon sunshine, common woolly sunflower

**var. *integritolium* (Hook.) Smiley [FNA21, HC, HC2]**

Univ. Calif. Publ. Bot. 9: 378.

Oregon sunshine

FNA21: "Variety *integritolium* intergrades with var. *lanatum* in Oregon and Washington near the Columbia River. The intermediate populations that have been analyzed are polyploid (J. S. Mooring 2001)."

**var. *lanatum* [FNA21, HC, HC2]**

Gen. Sp. Pl. 28.

common woolly sunflower

**var. *leucophyllum* (DC.) W.R. Carter [FNA21, HC2]**

Prelim. Cat. Fl. Vancouver. 82.

common woolly sunflower

***Eucephalus* [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 298. 1840.  
aster

***Eucephalus engelmannii* (D.C. Eaton) Greene [FNA20, HC2]**

Pittonia. 3: 54.

Engelmann's aster

*Aster engelmannii* (D.C. Eaton) A. Gray [HC]

***Eucephalus glaucescens* (A. Gray) Greene [FNA20, HC2]**

Pittonia. 3: 56.

Klickitat aster

*Aster engelmannii* (D.C. Eaton) A. Gray var. *glaucescens* A. Gray  
*Aster glaucescens* (A. Gray) S.F. Blake [HC]

FNA20: "Eucephalus glaucescens is known from the vicinity of Mt. Adams in Klickitat, Skamania, and Yakima counties. Intermediates with *E. ledophyllus* have been reported."

***Eucephalus ledophyllus* (A. Gray) Greene [FNA20, HC2]**

Pittonia. 3: 55.

Cascade aster

*Aster ledophyllus* (A. Gray) A. Gray [HC]

**var. *ledophyllus* [FNA20, HC2]**

Pittonia. 3: 55.

Cascade aster

*Aster ledophyllus* (A. Gray) A. Gray var. *ledophyllus* [HC]

***Eucephalus paucicapitatus* (B.L. Rob.) Greene [FNA20, HC2]**

Pittonia. 3: 56.

Olympic Mountain aster

*Aster paucicapitatus* (B.L. Rob.) B.L. Rob. [HC]

***Eurybia* [FNA20, HC2]**

Dict. Sci. Nat. ed. 2. 16: 46. 1820.

aster

***Eurybia conspicua* (Lindl.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 259.

western showy aster

*Aster conspicuus* Lindl. [HC]

***Eurybia integrifolia* (Nutt.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 260.

thickstem aster

*Aster amplexifolius* Rydb.

*Aster integrifolius* Nutt. [HC]

FNA20: "Eurybia integrifolia is found in mountain ranges bordering the Basin and Range Province, from the Sierra Nevada and Cascade ranges in the west to the Rocky Mountains and Colorado Plateau in the east."

***Eurybia merita* (A. Nelson) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 260.

subalpine aster

*Aster meritus* A. Nelson

*Aster sibiricus* L. ssp. *meritus* (A. Nelson) G.W. Douglas [ILBC1]

*Aster sibiricus* L. var. *meritus* (A. Nelson) Raup [HC]

FNA20: "Eurybia sibirica has often been confused at its southern range limit with *E. merita*, from which it differs by its often more low-cespitoso habit (versus more erect habit, but smaller individuals may be similar in this respect), usually more serrate leaves (versus subserrate to nearly entire), and subequal, foliaceous, purplish phyllaries (versus unequal, non-foliaceous, purple-margined). At the southern end of its range,

near the Canada?United States border, *E. sibirica* is usually found at higher elevations than its congener, there at its northern limit. *Aster sibiricus* forma *albinus* Lepage is merely a color variant of the species and is not recognized here."

***Eurybia radulina* (A. Gray) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 261.  
rough-leaved aster

*Aster radulinus* A. Gray [HC]

***Eurybia sibirica* (L.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 261.  
arctic aster

*Aster sibiricus* L. [HC]

*Aster sibiricus* L. ssp. *sibiricus*

FNA20: "Eurybia sibirica has often been confused at its southern range limit with *E. merita*, from which it differs by its often more low-cespitoso habit (versus more erect habit, but smaller individuals may be similar in this respect), usually more serrate leaves (versus subserrate to nearly entire), and subequal, foliaceous, purplish phyllaries (versus unequal, non-foliaceous, purple-margined). At the southern end of its range, near the Canada?United States border, *E. sibirica* is usually found at higher elevations than its congener, there at its northern limit. *Aster sibiricus* forma *albinus* Lepage is merely a color variant of the species and is not recognized here."

***Euthamia* [FNA20, HC2]**

Dict. Sci. Nat. ed. 2. 37: 471. 1825.  
grass-leaved goldenrod

***Euthamia graminifolia* (L.) Nutt. [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 325.  
fragrant goldenrod, grass-leaved goldenrod

*Euthamia graminifolia* (L.) Nutt. var. *major* (Michx.) Moldenke

*Solidago graminifolia* (L.) Salisb. [HC]

*Solidago graminifolia* (L.) Salisb. var. *major* (Michx.) Fernald [HC]

WA report in Abrams and FNA20, however no voucher at WTU.

***Euthamia occidentalis* Nutt. [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 326.  
western goldenrod, western goldentop

*Euthamia californica* Gand.

*Euthamia linearifolia* Gand.

*Solidago occidentalis* (Nutt.) Torr. & A. Gray [HC]

***Eutrochium* [FNA21, HC2]**

New Fl. 4: 78. 1838.  
Joe-pye weed

***Eutrochium maculatum* (L.) E.E. Lamont [FNA21, HC2]**

Sida. 21: 902.  
spotted Joe-pye weed

*Eupatoriadelphus maculatus* (L.) R.M. King & H. Rob.

*Eupatorium maculatum* L. [HC]

**var. *bruneri* (A. Gray) E.E. Lamont [FNA21, HC2]**

Sida. 21: 902. 2004.  
Joepyeweed, Joe-pye weed

*Eupatorium maculatum* L. ssp. *bruneri* (A. Gray) G.W. Douglas

*Eupatorium maculatum* L. var. *bruneri* (A. Gray) Breitung [HC]

(Douglas et al. 1998) state probably introduced in BC.

***Filago* [FNA19, HC, HC2]**

Sp. Pl. 2: 927, 1199. 1753; Gen. Pl. ed. 5, 397. 1754.  
filago

***Filago arvensis* L. [HC, HC2]**

Sp. Pl. 2: add.  
field cottonrose, field filago

*Logfia arvensis* (L.) Holub [FNA19]

*Oglifa arvensis* (L.) Cass.

FNA: "Logfia arvensis appears to be basal or nearly so in Logfia and Filagininae (J. D. Morefield 1992); only 2?4 epappose florets are present in most heads.....The earliest specimen confirmed from the flora area was from Bonner County, Idaho, in 1934."

***Filago vulgaris* Lam. [FNA19, HC2]**

Fl. Franç. 2: 61.  
German filago

*Filago germanica* L., homonym (illegitimate)

voucher? only WA report in Atkinson & Sharpe (1993) some BC records of this from the Gulf Islands have been annot. to *F. pyramidata*

***Gaillardia* [FNA21, HC, HC2]**

Observ. Phys. 29: 55. 1786 (as Gaillarda); Hist. Acad. Roy. Sci. Mém. Math. Phys. (Paris, 4to) 1786: 5. 1788.  
blanket-flower, gaillardia

***Gaillardia aristata* Pursh [FNA21, HC, HC2]**

Fl. Amer. Sept. 2: 573.  
blanket flower, great-flowered gaillardia

***Galinsoga* [FNA21, HC, HC2]**

Fl. Peruv. Prodr. 110, plate 24. 1794.  
garden pest, quickweed

**var. *parviflora* [FNA21, HC2]**

Icon. 3: 41, plate 281.  
small flowered galinsoga, gallant soldier

***Galinsoga quadriradiata* Ruiz & Pav. [FNA21, HC2]**

Syst. Veg. Fl. Peruv. Chil. 1: 198.  
ciliate galinsoga, shaggy galinsoga, quickweed

***Galinsoga ciliata* (Raf.) S.F. Blake [HC]**

***Gamochaeta* [FNA19, HC2]**

Chlor. Andina. 1: 151. 1856.  
cudweed

***Gamochaeta ustulata* (Nutt.) Holub [FNA19, HC2]**

Folia Geobot. Phytotax. 11: 83.  
purple cudweed, spoon-leaf cudweed

*Gamochaeta purpurea* (L.) Cabrera [FNA19], misapplied

*Gnaphalium purpureum* L. var. *ustulatum* (Nutt.) B. Boivin

FNA Volume 19:"Gamochaeta ustulata usually has been included in *G. purpurea*; it differs mostly in its longer duration, thicker and shorter stems, larger, more compact arrays of larger, brown heads, and aspects of phyllary morphology."

***Gnaphalium* [FNA19, HC, HC2]**

Sp. Pl. 2: 850. 1753; Gen. Pl. ed. 5, 368. 1754.  
cudweed, everlasting  
(see also *Gamochaeta*, *Pseudognaphalium*)

***Gnaphalium palustre* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 403.  
lowland cudweed, western marsh cudweed

*Filaginella palustris* (Nutt.) Holub  
*Gnaphalium heteroides* Klatt  
*Gnaphalium palustre* Nutt. var. *nanum* Jeps.

***Gnaphalium uliginosum* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 856.  
marsh cudweed

FNA19: "Gnaphalium uliginosum is native to Europe; it is not clear whether some or all of the North American plants may have been introduced into the flora."

***Grindelia* [FNA20, HC, HC2]**

Ges. Naturf. Freunde Berlin Mag. Neuesten Entdeck. Gesammten Naturk. 1: 259. 1807.  
grindelia, gumplant, gumweed, resinweed

***Grindelia integrifolia* DC. [FNA20, HC, HC2]**

Prodri. 5: 315.  
Puget Sound gumweed, Willamette Valley gumweed  
(see also *Grindelia hirsutula*)

*Grindelia integrifolia* DC. var. *integrifolia* [HC]

FNA20: "But for the stipitate-glandular apices of the phyllaries, plants of *Grindelia integrifolia* are very much like some plants treated here in *G. hirsutula*. Taxonomic status for plants that have been called *G. integrifolia* should be reconsidered."

***Grindelia nana* Nutt. [JPM2, OFP]**

Trans. Amer. Philos. Soc. ser. 2, 7: 314.  
Idaho gumweed, low gumweed

*Grindelia nana* Nutt. ssp. *nana*

var. *discoidea* (Nutt.) A. Gray [OFP]  
Syn. Fl. N. Amer. 1(2): 119.  
Columbia gumweed, rayless gumweed

*Grindelia columbiana* (Piper) Rydb. [HC]  
*Grindelia nana* Nutt. ssp. *columbiana* Piper

Endemic to the Columbia River Gorge in Washington and Oregon.

var. *nana* [Revised HC2, Flora of Oregon]  
low gumweed

*Grindelia hirsutula* Hook. & Arn. [FNA, HC2, OFP, JPM2], misapplied

***Grindelia squarrosa* (Pursh) Dunal [FNA20, HC, HC2]**

Mém. Mus. Hist. Nat. 5: 50.  
curlycup gumweed, serrate resinweed  
(see also *Grindelia hirsutula*)

*Grindelia squarrosa* (Pursh) Dunal var. *squarrosa* [HC]

Oregon Flora Project: "Although *G. squarrosa* is similar to *G. nana* in the form of its inflorescence and can sometimes be difficult to distinguish, the two species are rather distantly related and are unlikely to hybridize. *Grindelia squarrosa* appears to be increasing throughout its range and taking over areas that were occupied until recently by *G. nana* (and other species in other states)."

var. *serrulata* (Rydb.) Steyermark [OFP]  
curlycup gumweed, serrate resinweed  
(see also *Grindelia hirsutula*)

Oregon Flora Project: "Although *G. squarrosa* is similar to *G. nana* in the form of its inflorescence and can sometimes be difficult to distinguish, the two species are rather distantly related and are unlikely to

hybridize. *Grindelia squarrosa* appears to be increasing throughout its range and taking over areas that were occupied until recently by *G. nana* (and other species in other states)."

***Grindelia stricta* DC. [JPM2, OFP]**

Prod. [A. P. de Candolle] 7(1): 278. 1838.  
Oregon gumweed

<i>Grindelia hirsutula</i> is misapplied in Washington. Coastal specimens of this species are referential to <i>G. stricta</i> and <i>G. integrifolia</i>, and interior specimens are referential to <i>G. nana</i>.

**var. *stricta* [JPM2, OFP]**

*Grindelia stricta* DC. ssp. *stricta*

<i>Grindelia hirsutula</i> is misapplied in Washington. Coastal specimens of this species are referential to <i>G. integrifolia</i>, and interior specimens are referential to <i>G. nana</i>.

***Guizotia* [FNA21, HC2]**

Dict. Sci. Nat. ed. 2. 59: 237, 247, 248. 1829.

***Guizotia abyssinica* (L. f.) Cass. [FNA21, HC2]**

Dict. Sci. Nat. ed. 2. 59: 248. 1829.

*Polymnia abyssinica* L. f.

FNA21: "In the flora area, *Guizotia abyssinica* has been recorded sporadically at widely scattered stations (evidently often from birdseed wastes); it may be persistently established at relatively few stations."

***Gutierrezia* [FNA20, HC, HC2]**

Gen. Sp. Pl. 30. 1816.  
matchbrush, matchweed, snakeweed

***Gutierrezia sarothrae* (Pursh) Britton & Rusby [FNA20, HC, HC2]**

Trans. New York Acad. Sci. 7: 10. 1887.  
kindlingweed, matchweed, broom snakeweed

FNA20: "Gutierrezia sarothrae is often abundant in overgrazed pastures."

***Helenium* [FNA21, HC, HC2]**

Sp. Pl. 2: 886. 1753; Gen. Pl. ed. 5, 377. 1754.  
sneezeweed

***Helenium autumnale* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 886.  
common sneezeweed, large flowered sneezeweed

*Helenium autumnale* L. var. *autumnale*

*Helenium autumnale* L. var. *grandiflorum* Torr. & A. Gray [HC]  
*Helenium autumnale* L. var. *montanum* (Nutt.) Fernald [HC]

***Helianthella* [FNA21, HC, HC2]**

Fl. N. Amer. 2: 333. 1842.  
helianthella, little-sunflower

***Helianthella uniflora* (Nutt.) Torr. & A. Gray [FNA21, HC, HC2]**

Fl. N. Amer. 2: 334.  
Rocky Mountain helianthella  
var. *douglasii* (Torr. & A. Gray) W.A. Weber [FNA20, HC, HC2]  
Fl. N. Amer. 2: 334.  
Douglas helianthella, false sunflower

FNA20: "Two infraspecific taxa within *Helianthella uniflora* may be distinguished; they have been named at varietal rank. Variety *douglasii* has stems hirsute; involucres (15?)20?25(?30) mm diam.; outer phyllaries rarely elongated, margins ciliate, abaxial faces sparsely puberulent; ray laminae 30?40 mm; and 2n = 30. It grows in grasslands in the northern Rocky Mountains and on the east side of the Cascade Range (B.C.; Idaho, Oreg., Wash.) at 300?2500 m where it flowers May?Jul." See Weber

(1952)

***Helianthus* [FNA21, HC, HC2]**

Sp. Pl. 2: 904. 1753; Gen. Pl. ed. 5, 386. 1754.  
sunflower

***Helianthus annuus* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 904.  
common sunflower

*Helianthus annuus* L. ssp. *lenticularis* (Douglas ex Lindl.) Cockerell

FNA21: "Helianthus annuus is widely distributed, including weedy, cultivated, and escaped plants. It is the only native North American species to become a major agronomic crop. Despite its considerable variability, attempts have failed to produce a widely adopted infraspecific system of classification. Forms with red-colored ray laminae, known from cultivation and occasionally seen escaped, trace their ancestry to a single original mutant plant. It hybridizes with many of the other annual species."

***Helianthus ciliaris* DC. [FNA21, HC2]**

Prod. 5: 587.  
blueweed

***Helianthus cusickii* A. Gray [FNA21, HC, HC2]**

Proc. Amer. Acad. Arts. 21: 413.  
Cusick's sunflower, turniproot sunflower

***Helianthus nuttallii* Torr. & A. Gray [FNA21, HC, HC2]**

Fl. N. Amer. 2: 324. 1842.  
Nuttall's sunflower

**ssp. *nuttallii* [FNA21, HC2]**

Fl. N. Amer. 2: 324.  
cordilleran sunflower, Nuttall's sunflower

*Helianthus nuttallii* Torr. & A. Gray var. *nuttallii* [HC]

***Helianthus petiolaris* Nutt. [FNA21, HC, HC2]**

J. Acad. Nat. Sci. Philadelphia. 2: 115.  
prairie sunflower

*Helianthus petiolaris* Nutt. var. *petiolaris*

**ssp. *petiolaris* [FNA21, HC2]**  
prairie sunflower

***Helianthus tuberosus* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 905.  
Jerusalem artichoke

***Hemizonella* [FNA21, HC2]**

Proc. Amer. Acad. Arts. 9: 189. 1874.  
miniature tarweed

***Hemizonella minima* (A. Gray) A. Gray [FNA21, HC2]**

Proc. Amer. Acad. Arts. 9: 189.  
smallhead tarplant, least tarweed, small-head tarweed, smallhead tarweed

*Hemizonia minima* A. Gray

*Madia minima* (A. Gray) D.D. Keck [HC]

FNA21: "Hemizonella minima is self-compatible, like most other tarweeds that are distributed widely in western North America."

***Heterotheca* [FNA20, HC2]**

Bull. Sci. Soc. Philom. Paris. 1817: 137. 1817.  
camphorweed, goldaster

***Heterotheca oregona* (Nutt.) Shinners [FNA20, HC2, OFP]**

Field & Lab. 19: 71.

Oregon goldenaster

*Chrysopsis oregona* (Nutt.) A. Gray [HC]

*Chrysopsis oregona* (Nutt.) A. Gray var. *oregona*

var. ***oregona*** [FNA20, HC2]

Oregon goldenaster

***Heterotheca villosa* (Pursh) Shinners [FNA20, HC2]**

Field & Lab. 19: 71. 1951.

hairy goldaster

*Chrysopsis villosa* (Pursh) Nutt. ex DC. [HC]

var. ***foliosa*** (Nutt.) V.L. Harms [FNA20, HC2]

Wrightia. 4: 15. 1968.

leafy goldenaster

*Chrysopsis villosa* (Pursh) Nutt. ex DC. var. *foliosa* (Nutt.) Cronquist [HC]

*Chrysopsis villosa* (Pursh) Nutt. ex DC. var. *foliosa* (Nutt.) D.C. Eaton, orthographic variant

Voucher? OR reports in Peck (1961) misapplied (Chambers and Sundberg 2000). FNA20: "Variety *foliosa* is most common at the base of the Front Range in Colorado and Wyoming; it occurs scattered across the northern part of its range in the mountains and western prairies. Plants that are glandular but otherwise like var. *foliosa* are presumed here to be hybrids with either var. *minor* or var. *nana*. Variety *foliosa* is most similar to var. *ballardii*."

var. ***minor*** (Hook.) Semple [FNA20, HC2]

Novon. 4: 54. 1994.

hairy goldenaster, hispid goldenaster

*Chrysopsis villosa* (Pursh) Nutt. ex DC. var. *hispida* (Hook.) A. Gray [HC]

*Heterotheca villosa* (Pursh) Shinners var. *hispida* (Hook.) V.L. Harms

FNA20: "Variety *minor* is distinguished by its usually narrowly to broadly oblanceolate distal cauline leaves (sometime oblong or ovate) and moderately glandular and hispido-strigose indument. The leaf bases are usually narrowly to broadly cuneate or attenuate (rarely rounded). Distal leaf faces usually have about 10?50 hairs/mm<sup>2</sup> and about 4?20 glands/mm<sup>2</sup>. The variety is the most variable in the species and includes some local distinctive morphotypes that grade into other forms. Plants intermediate between this and all other varieties occur in areas where the ranges are sympatric, and they make infraspecific taxonomy of the species difficult. The variety has been incorrectly referred to as var. *hispida* (a later synonym) in most floras. The status of *Heterotheca barbata* (Rydberg) Semple (*Chrysopsis barbata* Rydberg), the Spokane goldenaster, is uncertain. J. C. Semple (1996) treated it as a separate species to draw attention to the problem; a detailed description based on the type and detailed illustrations were included. It is known from the type collection along the Spokane River Valley east of Spokane, Idaho, and two down-river, atypical collections (Benton and Spokane counties, Washington). It flowers in July (sometimes August). It is similar to *H. villosa* var. *minor*, but differs in having lanceolate-elliptic distal cauline leaves (34?38 × 8?9 mm) that are little reduced distally, long branches (each with one to a few large heads), and disc corollas with a few, very long hairs on the tube. Further work is needed to increase the number of specimens available for a detailed comparison with *H. villosa* var. *minor* in order to clarify whether *H. barbata* warrants species level recognition, should be included in *H. villosa* as a variety, or placed in synonymy under var. *minor*."

var. ***villosa*** [FNA20, HC2]

Field & Lab. 19: 71.

hairy goldaster, hairy goldenaster

*Chrysopsis villosa* (Pursh) Nutt. ex DC. var. *villosa* [HC]

***Hieracium* [FNA19, HC, HC2]**

Sp. Pl. 2: 799. 1753; Gen. Pl. ed. 5, 350. 1754.

hawkweed

***Hieracium albiflorum*** Hook. [FNA19, HC, HC2]

Fl. Bor.-Amer. 1: 298.  
white hawkweed, white-flowered hawkweed

*Hieracium aurantiacum* L. [FNA19, HC, HC2]  
Sp. Pl. 2: 801.  
orange hawkweed

*Hieracium caespitosum* Dumort. [FNA19, HC2]  
Fl. Belg. 62.  
meadow hawkweed, yellow king devil

*Hieracium pratense* Tausch [HC]

*Hieracium flagellare* Willd. [FNA19, HC2]  
Enum. Pl. suppl.: 54.  
whip hawkweed

*Hieracium flagellare* Willd. var. *amauracron* (Missback & Zahn) Lepage [KZ99]  
*Hieracium flagellare* Willd. var. *cernuiforme* (Naegeli & Peter) Lepage [KZ99]  
*Hieracium ×flagellare* Willd. var. *flagellare* [KZ99]  
*Hieracium flagellare* Willd. var. *pilosus* Lepage [KZ99]

FNA19: "The type of *Hieracium flagellare* may have resulted from a cross between plants of *H. caespitosum* and *H. pilosella* (A. Cronquist 1980)." Although generally treated as a hybrid, this plant is often found in the absence of both of its parents, and is a widespread weed in eastern North America (Lepage 1967). Recently collected in San Juan County.

*Hieracium ×floribundum* Wimm. & Grab. [HC2]  
flowery hawkweed

Also written as *Hieracium floribundum*.

*Hieracium glomeratum* Froel. [HC2]  
Prodromus Systematis Naturalis Regni Vegetabilis 7(1): 207.  
yellow devil hawkweed

*Hieracium lachenalii* C.C. Gmel. [HC2]  
Fl. Bad. 3: 322.  
common hawkweed, English hawkweed, European hawkweed

*Hieracium vulgatum* Fr. [FNA19, HC], misapplied  
FNA19: "The correct name for the species here called *Hieracium vulgatum* may be *H. lachenallii* C. C. Gmelin (E. Lepage 1971; E. G. Voss 1972?1996, vol. 3)." Weedy on the west side of the Cascades (Zika 2002), where first collected in 1966.

*Hieracium longiberbe* Howell [FNA19, HC, HC2]  
Fl. N.W. Amer. 395.  
long-bearded hawkweed

FNA19: "*Hieracium longiberbe* is known only from along the Columbia River."

*Hieracium maculatum* Sm. [HC2, Stace 1997]  
Baier. Fl. 2: 319.  
mottled hawkweed

*Hieracium murorum* L. [FNA19, HC2]  
Sp. Pl. 2: 802.  
wall hawkweed

*Hieracium pilosella* L. [FNA19, HC, HC2]  
Sp. Pl. 2: 800.  
mouse-ear hawkweed  
*Hieracium pilosella* L. var. *nivea* Muell.Arg. [KZ99]  
*Hieracium pilosella* L. var. *pilosella* [KZ99]  
*Pilosella officinarum* F.W. Schultz & Sch. Bip.

*Hieracium piloselloides* Vill. [FNA19, HC2]

Prosp. Hist. Pl. Dauphiné. 34.

tall hawkweed

*Hieracium florentinum* All.

FNA19: "Plants called *Hieracium praealtum* Villars ex Gochnat (at least those called *H. praealtum* var. *decipliens* W. D. J. Koch) reputedly differ from members of *H. piloselloides* in having blades of their proximal leaves stellate-pubescent abaxially (M. L. Fernald 1950); such plants may be found in the flora and may merit taxonomic recognition."

***Hieracium sabaudum* L. [FNA19, HC2]**

Sp. Pl. 2: 804.

savoy hawkweed

*Hieracium laevigatum* Willd., misapplied

***Hieracium scouleri* Hook. [FNA19, HC, HC2]**

Fl. Bor.-Amer. 1: 298.

hound-tongue hawkweed, Scouler's hawkweed

*Hieracium albertinum* Farr [HC]

*Hieracium chapacanum* Zahn

*Hieracium cusickii* Gand.

*Hieracium cynoglossoides* Arv.-Touv. [HC]

*Hieracium scouleri* Hook. var. *albertinum* (Farr) G.W. Douglas & G.A. Allen

*Hieracium scouleri* Hook. var. *griseum* A. Nelson

*Hieracium scouleri* Hook. var. *scouleri*

***Hieracium ×stoloniflorum* Waldst. & Kit. [HC2]**

Descr. Icon. Pl. Hung. 3: 303, t. 273.

forked hawkweed

(= *Hieracium aurantiacum* × *Hieracium pilosella*)

*Hieracium ×stoloniflorum* Waldst. & Kit. var. *cayouetteanum* Lepage [KZ99]

*Hieracium ×stoloniflorum* Waldst. & Kit. var. *laurentianum* Lepage [KZ99]

*Hieracium ×stoloniflorum* Waldst. & Kit. var. *stoloniflorum* [KZ99]

***Hieracium triste* Willd. ex Spreng. [FNA19, HC2]**

Syst. Veg. 3: 640.

alpine hawkweed, slender hawkweed

*Hieracium gracile* Hook. [HC]

*Hieracium gracile* Hook. var. *densifloccosum* (Zahn) Cronquist

*Hieracium gracile* Hook. var. *detonsum* (A. Gray) A. Gray

*Hieracium gracile* Hook. var. *gracile*

*Hieracium triste* Willd. ex Spreng. var. *gracile* (Hook.) A. Gray

***Hieracium umbellatum* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 804.

narrowleaf hawkweed, umbellate hawkweed

*Hieracium canadense* Michx. [HC]

*Hieracium kalmii* L.

*Hieracium umbellatum* L. ssp. *umbellatum*

FNA19: "The circumscription of *Hieracium umbellatum* adopted here is supported by research done by others, especially G. A. Guppy (1978) and E. Lepage (1960). *Hieracium canadense* var. *kalmii* (Linnaeus) Scoggan, referable here, is an illegitimate name."

***Hulsea* [FNA21, HC, HC2]**

Pacif. Railr. Rep. 6(3): 77, plate 13. 1858.

alpinegold, hulsea

***Hulsea nana* A. Gray [FNA21, HC, HC2]**

Pacif. Railr. Rep. 6(3): 76, plate 13.

small alpinegold, small hulsea

*Hulsea nana* A. Gray var. *larsenii* A. Gray

FNA19: "Densely lanate or woolly plants of *Hulsea nana* are referable to var. *larsenii*. Such plants may occur in distinct populations but can be found together with sparsely lanate and strictly glandular plants. The distribution of lanate to woolly plants appears associated with higher levels of insolation."

***Hymenopappus* [FNA21, HC, HC2]**

Hymenopappus. plate. 1788.

hymenopappus

***Hymenopappus filifolius* Hook. [FNA21, HC, HC2]**

Fl. Bor.-Amer. 1: 317.

Columbia cut-leaf, fineleaf hymenopappus

**var. *filifolius* [FNA21, HC, HC2]**

Columbia cutleaf

***Hypochaeris* [FNA19, HC, HC2]**

Sp. Pl. 2: 810. 1753; Gen. Pl. ed. 5, 352. 1754.

cats-ear

***Hypochaeris glabra* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 811.

smooth cat's-ear

FNA19: "*Hypochaeris glabra* is usually distinguishable by its annual habit and relatively small size, slender and shallow roots, fine stems, often glabrous leaves, and beakless, truncate outer cypselae. Occasional specimens are larger and have induments characteristics of *H. radicata*; they can be distinguished by the dimorphic cypselae."

***Hypochaeris radicata* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 811.

hairy cat's-ear, rough cat's-ear

FNA19: "<i>Hypochaeris radicata</i> is recognized by the coarse, perennial habit, stout roots, coarsely hirsute leaves and phyllaries, yellow corollas, and monomorphic, beaked cypselae. It is weedy and invasive in some areas."

***Inula* [FNA19, HC, HC2]**

Sp. Pl. 2: 881. 1753; Gen. Pl. ed. 5, 375. 1754.

inula

***Inula helenium* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 881.

elecampane, inula

Known in WA from a single collection in Stevens County.

***Ionactis* [FNA20, HC2]**

Pittonia. 3: 245. 1897.

ankle-aster

***Ionactis stenomeres* (A. Gray) Greene [FNA20, HC2]**

Pittonia. 3: 246.

Rocky Mountain ankle-aster, Rocky Mountain aster

Aster *stenomeres* A. Gray [HC]

***Iva* [FNA21, HC, HC2]**

Sp. Pl. 2: 988. 1753; Gen. Pl. ed. 5, 426. 1754.

poverty-weed

(see also *Cyclachaena*)

***Iva axillaris* Pursh [FNA21, HC, HC2]**

Fl. Amer. Sept. 2: 743.

deeproot, poverty weed

*Iva axillaris* Pursh ssp. *robustior* (Hook.) Bassett

***Jacobaea* [HC2]**

ragwort

***Jacobaea maritima* (L.) Pelser & Meijden [HC2]**

Heukels' Fl. Nederland 677.

silver ragwort

***Jacobaea vulgaris* Gaertn. [HC2]**

Fruct. Sem. Pl. 2: 445.

tansy ragwort

***Senecio jacobaea* L. [FNA20, HC]**

FNA20: "Senecio jacobaea is a weed introduced from Europe and now well established in places of cool, damp summers. It is toxic to livestock and legally noxious in most states and provinces where it occurs. The Russian botanist E. Wiebe (2000) resuscitated Jacobaea for plants that are treated here as Senecio jacobaea, S. erucifolius, and S. cannabifolius. Phylogenetic studies may confirm the utility of recognizing Jacobaea as a distinct genus; to do so here would be premature."

***Jaumea* [FNA21, HC, HC2]**

Syn. Pl. 2: 397. 1807.

jaumea

***Jaumea carnosa* (Less.) A. Gray [FNA21, HC, HC2]**

U.S. Expl. Exped. 17: 360.

fleshy jaumea, marsh jaumea

*Coinogyne carnosa* Less.

***Lactuca* [FNA19, HC, HC2]**

Sp. Pl. 2: 795. 1753; Gen. Pl. ed. 5, 348. 1754.

lettuce

(see also *Mycelis*)

***Mulgedium* [FNA19]**

***Lactuca biennis* (Moench) Fernald [FNA19, HC, HC2]**

Rhodora. 42: 300.

tall blue lettuce, wild blue lettuce

*Sonchus biennis* Moench

FNA19: "The type of *Lactuca terrae-novae* Fernald is probably conspecific with that of *L. biennis*. The type of *L. biennis* may be conspecific with that of *L. floridana*."

***Lactuca canadensis* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 796.

Canadian wild lettuce, Florida blue lettuce

***Lactuca ludoviciana* (Nutt.) Riddell [FNA19, HC, HC2]**

W. J. Med. Phys. Sci. 8: 491.

Louisiana lettuce, prairie lettuce, western lettuce

***Lactuca saligna* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 796.

least lettuce, willow lettuce

***Lactuca sativa* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 795.

garden lettuce

***Lactuca serriola* L. [FNA19, HC, HC2]**

Cent. Pl. II. 29.

prickly lettuce

*Lactuca scariola* L. var. *integrata* Gren. & Godr.  
*Lactuca scariola* L. var. *scariola*

***Lactuca tatarica* (L.) C.A. Mey. [HC2]**

Verz. Pfl. Casp. Meer. (C.A. von Meyer). 56.  
blue lettuce

**ssp. *pulchella* (Pursh) Stebbins [HC2]**

Madroño 5: 123.  
blue lettuce

*Lactuca pulchella* (Pursh) DC. [HC]

*Lactuca tatarica* (L.) C.A. Mey. var. *pulchella* (Pursh) Breitung  
*Mulgedium oblongifolium* (Nutt.) Reveal  
*Mulgedium pulchellum* (Pursh) G. Don [FNA19]

FNA19: "The type of *Mulgedium pulchellum* may be conspecific with that of *M. tataricum* (Linnaeus) de Candolle, a Eurasian species. Or, if "perennial" plus "Fl. blue" constitutes sufficient description for valid publication of the name *Lactuca oblongifolia* Nuttall (1813), then a new combination in *Mulgedium* based on that name may be appropriate for what is here called *M. pulchellum*."

***Lactuca virosa* L. [FNA19, HC2]**

Sp. Pl. 2: 795.  
great lettuce, tall lettuce, wild lettuce

***Lagophylla* [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 390. 1841.  
hareleaf, rabbitleaf

***Lagophylla ramosissima* Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 391.  
slender hareleaf, common rabbitleaf

*Lagophylla ramosissima* Nutt. ssp. *ramosissima*

FNA21: "*Lagophylla ramosissima* occurs widely in dry, often disturbed or poor soils of the California Floristic Province, Great Basin, and Pacific Northwest. Plants with heads in glomerate arrays have been treated as *L. congesta* or *L. ramosissima* subsp. *congesta*; W. C. Thompson (1983, p. 21) concluded that *L. congesta* represents an "extreme morphological variant of *L. ramosissima*" unworthy of taxonomic recognition." Chambers and Sundberg (2000) question if ssp. *congesta* (Greene) D.D. Keck in CA is a good taxon

***Lapsana* [FNA19, HC, HC2]**

Sp. Pl. 2: 811. 1753; Gen. Pl. ed. 5, 353. 1754.  
nipplewort

***Lapsana communis* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 811. 1753.  
common nipplewort

FNA19: "<i>Lapsana communis</i> is widely distributed in North America. It is easily recognized by the abruptly constricted lyrate leaves with relatively large terminal lobes, heads of relatively small flowers with yellow corollas, keeled phyllaries, and epappose cypselae. It is aggressively weedy and often found in shady disturbed sites. The milky juice of <i>L. communis</i> is said to be soothing to sensitive skin, particularly on the nipples of nursing mothers."

***Lasthenia* [FNA21, HC, HC2]**

Opusc. Phytol. 3: 88. 1834.  
goldfields

***Lasthenia glaberrima* DC. [FNA21, HC, HC2]**

Prodri. 5: 664.  
smooth goldfields, smooth lasthenia

***Lasthenia maritima* (A. Gray) M.C. Vasey [FNA21, HC2]**

Madroño. 32: 139.

maritime goldfields, seaside goldfields

*Baeria maritima* (A. Gray) A. Gray

*Baeria minor* (DC.) Ferris ssp. *maritima* (A. Gray) Ferris

*Lasthenia minor* (DC.) Ornduff var. *maritima* (A. Gray) Cronquist [HC]

FNA21: "Lasthenia maritima is a self-pollinating, "guano endemic" of seabird nesting grounds. It is typically found on offshore islands and rocks from the Farallon Islands, California, to the northern tip of Vancouver Island, British Columbia and rarely occurs on the mainland."

***Lasthenia minor* (DC.) Ornduff [FNA21, HC, HC2]**

Univ. Calif. Publ. Bot. 40: 80.

coastal goldfields

*Baeria minor* (DC.) Ferris

FNA21 lists this as endemic to California, however Eugene Kozloff collected a plant in 1990 annotated to this name by Robert Ornduff. This determination was confirmed by D. Giblin in 2008 after comparison with *L. maritima* and *L. minor* specimens.

***Layia* [FNA21, HC, HC2]**

Prodr. 7: 294. 1838.

layia, tidytips

***Layia glandulosa* (Hook.) Hook. & Arn. [FNA21, HC, HC2]**

Bot. Beechey Voy. 358.

white layia, white daisy tidytips

*Layia glandulosa* (Hook.) Hook. & Arn. ssp. *glandulosa*

*Layia glandulosa* (Hook.) Hook. & Arn. ssp. *lutea* D.D. Keck

FNA21: "Layia glandulosa occurs in deserts of western North America, extending to the Pacific coast in central and southern California. As treated here (provisionally) and previously, *L. glandulosa* corresponds to a paraphyletic group; molecular phylogenetic data have indicated that *L. discoidea* is most closely related to a subset of lineages in *L. glandulosa*, including yellow-rayed populations previously recognized as subsp. *lutea* or var. *lutea* (B. G. Baldwin, unpubl.). Report of *L. glandulosa* from British Columbia has not been confirmed."

***Leontodon* [FNA19, HC, HC2]**

Sp. Pl. 2: 798. 1753; Gen. Pl. ed. 5, 349. 1754.

hawkbit

***Leontodon autumnalis* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 798. 1753.

autumn hawkbit

*Leontodon autumnalis* L. ssp. *autumnalis*

FNA19: "Leontodon autumnalis is recognized by the usually branched stems with (1)2-5 heads, peduncles bracteate proximal to heads, non-beaked cypselae, and pappi wholly of plumose bristles. It is now established in eastern North America and is sporadic in the west. Specimens with coarsely hirsute phyllaries have been recognized as var. *pratensis*; intermediates occur and the characteristic does not seem to correlate with other characters." Stace (1997) notes distinctness of subspp. & vars. need study

***Leontodon saxatilis* Lam. [FNA19, HC2]**

Fl. Franç. 2: 115.

hairy hawkbit

**ssp. *saxatilis* [FNA19, HC2]**

Fl. Franç. 2: 115.

lesser hawkbit

*Leontodon leysseri* (Wallr.) G. Beck

*Leontodon nudicaulis* Mérat [HC]

*Leontodon nudicaulis* Mérat ssp. *taraxacoides* (Vill.) Schinz & Thell. [HC]

*Leontodon taraxacoides* (Vill.) Mérat ssp. *taraxacoides*

*Lomatium nudicaulis* Mérat

***Leucanthemum* [FNA19, HC2]**

Gard. Dict. Abr. ed. 4. vol. 2. 1754.

daisy

***Leucanthemum ×superbum* (Bergmans ex J.W. Ingram) D.H. Kent [FNA19, HC2]**

Watsonia 18(1): 89.

Shasta daisy

voucher? WA report A. Jacobson (pers. comm.) Seattle area reseeding in gardens, truly naturalized? FNA19: "The name Shasta daisy of horticulture is associated also with *Leucanthemum ×superbum* (Bergmans ex J. Ingram) Bergmans ex D. H. Kent, which is generally thought to have been derived from hybrids between *L. maximum* and *L. lacustre*. Cultivars of "Shasta daisy" number in the dozens, including "single," "double," "quill," and "shaggy" forms; they may be encountered as waifs or persisting from abandoned plantings."

***Leucanthemum vulgare* Lam. [FNA19, HC2]**

Fl. Franç. 2: 137.

oxeye daisy

***Chrysanthemum leucanthemum* L. [HC]**

*Chrysanthemum leucanthemum* L. var. *pinnatifidum* Lecoq & Lamotte

***Logfia* [FNA19, HC2]**

Bull. Sci. Soc. Philom. Paris. 1819: 143. 1819 (as "genre ou sous-genre"). in F. Cuvier, Dict. Sci. Nat. ed. 2, 23: 564. 1822.

cottonrose, cottonweed

***Logfia gallica* (L.) Cosson & Germain [FNA19, HC2]**

Ann. Sci. Nat., Bot., sér. 2. 20: 291.

daggerleaf cottonweed

***Logfia minima* (Sm.) Dumort. [FNA19, HC2]**

Fl. Belg. 68.

little cottonrose

*Filago minima* (Sm.) Pers.

***Luina* [FNA20, HC, HC2]**

Hooker? s Icon. Pl. 12: 35, plate 1139. 1873.

luina

(see also *Cacaliopsis*, *Rainiera*)

***Luina hypoleuca* Benth. [FNA20, HC, HC2]**

Hooker? s Icon. Pl. 12: 36, plate 1139.

littleleaf luina, silverback luina, littleleaf silverback

***Lygodesmia* [FNA19, HC, HC2]**

Edinburgh New Philos. J. 6: 311. 1829.

rush-pink, skeletonplant

(see also *Pleiacanthus*)

***Lygodesmia juncea* (Pursh) D. Don ex Hook. [FNA19, HC, HC2]**

Fl. Bor.-Amer. 1: 295.

rush skeletonplant

FNA19: "Lygodesmia juncea is the most widespread species of the genus, occurring throughout the High Plains region of North America. It is easily distinguished by its bushy habit, greatly reduced cauline leaves, relatively small heads and involucres, and phyllaries lacking appendages. Mature cypselae are rarely found on this species, and the plants are presumably sterile and reproduce mainly by vegetative means. Many specimens have round galls to 10 mm diameter on the stems, produced by solitary wasps and apparently

unique to this species."

***Madia* [FNA21, HC, HC2]**

Sag. Stor. Nat. Chili. 136, 354. 1782.

madia, tarweed

(see also *Anisocarpus*, *Hemizonella*)

***Madia citriodora* Greene [FNA21, HC, HC2]**

Bull. Torrey Bot. Club. 9: 63.

lemon-scented tarplant, lemon tarweed, lemon-scented tarweed

***Madia elegans* D. Don ex Lindl. [FNA21, HC, HC2]**

Edwards's Bot. Reg. 17: plate 1458.

common madia, autumn showy tarweed

*Madia elegans* D. Don ex Lindl. ssp. *densiflora* (Greene) D.D. Keck

*Madia elegans* D. Don ex Lindl. ssp. *elegans*

*Madia elegans* D. Don ex Lindl. ssp. *vernalis* D.D. Keck

*Madia elegans* D. Don ex Lindl. var. *densifolia* (Greene) Jeps. [HC]

*Madia elegans* D. Don ex Lindl. var. *elegans* [HC]

FNA21: "Madia elegans occurs widely in California outside the deserts and in southwestern Oregon and locally in western Nevada and Washington. It is unusually variable in morphology, ecology, and phenology. Molecular data have indicated that D. D. Keck's (1959) infraspecific taxonomy for *M. elegans* needs revision."

***Madia exigua* (Sm.) A. Gray [FNA21, HC, HC2]**

Proc. Amer. Acad. Arts. 8: 391.

threadstem madia, little tarplant, little tarweed

FNA21: "Madia exigua occurs in seasonally dry situations in much of western North America outside the warm deserts. Morphologically, *M. exigua* is somewhat similar to *Hemizonella minima*, which (unlike *M. exigua*) has subumbellate arrays of heads and obcompressed, sparsely hairy ray cypselae."

***Madia glomerata* Hook. [FNA21, HC, HC2]**

Fl. Bor.-Amer. 2: 24.

mountain tarplant, cluster tarweed, mountain tarweed

***Madia gracilis* (Sm.) D.D. Keck [FNA21, HC, HC2]**

Madroño. 5: 169.

grassy tarplant, common tarweed, slender tarweed

*Madia gracilis* (Sm.) D.D. Keck ssp. *gracilis*

Hybridizes with *M. citriodora*, *M. sativa*

***Madia sativa* Molina [FNA21, HC, HC2]**

Sag. Stor. Nat. Chili. 136.

Chilean tarplant, coast tarweed

*Madia capitata* Nutt.

*Madia sativa* Molina ssp. *capitata* (Nutt.) Piper

*Madia sativa* Molina ssp. *sativa*

*Madia sativa* Molina var. *congesta* Torr. & A. Gray [HC]

*Madia sativa* Molina var. *sativa* [HC]

***Matricaria* [FNA19, HC, HC2]**

Sp. Pl. 2: 890. 1753; Gen. Pl. ed. 5, 380. 1754.

chamomile, matricaria, mayweed

(see also *Tripleurospermum*)

***Matricaria chamomilla* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 891.

wild chamomile, scented mayweed

*Matricaria recutita* L.

FNA19: "Although the name *Matricaria chamomilla* has been considered to be misapplied (e.g., S. Rauschert 1974; A. Cronquist 1994; E. G. Voss 1972?1996, vol. 3), W. L. Applequist (2002) argued convincingly that the name is indeed correctly applied to the taxon described here. Among the North American material, specimens with coronate ray cypselae (var. *chamomilla*), or wholly without coronas [var. *recutita* (Linnaeus) Grierson] have been encountered but none with fully coronate cypselae (var. *coronata* J. Gay ex Boissier), even though synonymy under this name includes *M. currantiana*, reported for Texas and New Mexico (specimens not seen). The varieties may not be worth recognizing (Applequist; Q. O. N. Kay 1976) and are not treated formally here." Rauschert (1974) transferred this from *Matricaria* to *Chamomilla*

***Matricaria discoidea* DC. [FNA19, HC2]**

Prod. 6: 50.

pineapple weed

*Chamomilla discoidea* (DC.) J. Gay ex A. Braun

*Chamomilla suaveolens* (Pursh) Rydb.

*Matricaria matricarioides* (Less.) Porter [HC], illegitimate name

*Santolina suaveolens* Pursh

FNA19: "Matricaria discoidea has been used as a medicinal and aromatic plant by Native American tribes (D. E. Moerman 1998). It also is considered a weed, and it is resistant to a photosystem II inhibitor herbicide in the United Kingdom ([www.weedscience.org](http://www.weedscience.org)). It is a northwestern North American native that has spread to eastern and northern North America and elsewhere (E. McClintock 1993b; E. G. Voss 1972?1996, vol. 3; A. Cronquist 1994). NatureServe ([www.natureserve.org](http://www.natureserve.org)) and Natural Resources Conservation Service ([plants.usda.gov](http://plants.usda.gov)) erroneously present *M. discoidea* as introduced on the continent. Its natural habitat is ill-defined because the species has become ruderal even in its native range. For discussion of the nomenclature of this taxon, see S. Rauschert (1974); K. N. Gandhi and R. D. Thomas (1991); Cronquist; and Voss. *Matricaria matricarioides* (Lessing) Porter cannot be applied to the American taxon; *M. matricarioides* was originally published as *Artemisia matricarioides* Lessing, a new name for *Tanacetum pauciflorum* Richardson (see S. Rauschert 1974), itself a synonym of *T. huronense* Nuttall. W. Greuter (pers. comm.), who accepts *M. discoidea*, considers Rauschert's treating *Artemisia matricarioides* as homotypic with *T. pauciflorum* as equivalent to a lectotype designation."

***Mauranthemum* [FNA19, HC2]**

Taxon. 44: 377. 1995.

***Micropus californicus* Fisch. & C.A. Mey. [FNA19, HC, HC2]**

Index Seminum (St. Petersburg). 2: 42.

cottontop, Q-tips

**var. *californicus* [FNA19, HC, HC2]**

cottontop, Q-tips

Photographed in Klickitat County by R.L. Carr in 2015:  
<http://web.ewu.edu/ewflora/Asteraceae/Micropus%20californicus.html>

***Microseris* [FNA19, HC, HC2]**

Philos. Mag. Ann. Chem. 11: 388. 1832.

microseris

(see also *Nothocalais*, *Uropappus*)

***Apargidium* [HC]**

***Microseris bigelovii* (A. Gray) Sch. Bip. [FNA19, HC, HC2]**

Jahresber. Pollichia. 22?24: 308.

coast microseris, coastal silverpuffs

FNA19: "Microseris bigelovii is the most characteristically coastal of the annual taxa and the only one to include plants with obtuse, spatulate leaves (K. Bachmann et al. 1984). A statistical analysis of its morphologic variation was published by Bachmann (1992). It sometimes has been collected at inland sites at 500?600 m, where the cypselae may have been introduced by domestic animals. The northern populations near Victoria, British Columbia, and the San Juan Islands, Washington, are disjunct from the main range, which extends from Oregon to Santa Barbara County, California."

***Microseris borealis* (Bong.) Sch. Bip. [FNA19, HC2]**

Jahresber. Pollichia. 22?24: 310.  
apargidium, bog microseris, northern silverpuffs

*Apargia borealis* Bong.

*Apargidium boreale* (Bong.) Torr. & A. Gray [HC]  
*Scorzonella borealis* (Bong.) Greene

***Microseris laciniata* (Hook.) Sch. Bip. [FNA19, HC, HC2]**

Jahresber. Pollichia. 22?24: 309.  
cut-leaved microseris

**ssp. *laciniata* [FNA19, HC2]**

Jahresber. Pollichia. 22?24: 309.  
cutleaf microseris, cut leaved scorzonella

*Scorzonella laciniata* (Hook.) Sch. Bip. var. *laciniata*

*Scorzonella laciniata* (Hook.) Sch. Bip. var. *pratensis* (Greene) Jeps.

*Scorzonella procera* (A. Gray) Greene

FNA19: "Subspecies laciniata occurs principally away from the coast, in interior valleys and hills, rarely reaching high elevations. The width of the outer phyllaries is a convenient way to separate it from subsp. leptosepala, with which it intergrades in the Klamath Mountains and at various sites east of the Cascade Range."

**ssp. *leptosepala* (Nutt.) K.L. Chambers [FNA19, HC2]**

Contr. Dudley Herb. 5: 61.  
cut-leaved silverpuffs, cutleaf silverpuffs

*Microseris leptosepala* (Nutt.) A. Gray

FNA19: "Subspecies leptosepala is known from the Klamath Mountains of California and Oregon and rare northward."

***Microseris nutans* (Hook.) Sch. Bip. [FNA19, HC, HC2]**

Jahresber. Pollichia. 22?24: 309. 1866.  
nodding microseris, nodding scorzonella, nodding silverpuffs

*Microseris nutans* (Hook.) Sch. Bip. ssp. *nutans*

*Scorzonella nutans* Hook.

*Scorzonella nutans* Hook. var. *major* (A. Gray) M. Peck

***Mycelis* [FNA19, HC2]**

Dict. Sci. Nat. ed. 2. 33: 483. 1824.  
mycelis

***Mycelis muralis* (L.) Dumort. [FNA19, HC2]**

Fl. Belg. 60.  
wall lettuce

*Lactuca muralis* (L.) Gaertn. [HC]

***Nabalus* [HC2]**

nabalus, rattlesnake-root

***Nabalus hastatus* (Less.) A.Heller [WTU]**

Muhlenbergia 1: 8.  
western white lettuce, western rattlesnake-root

*Nabalus alatus* Hook.

*Prenanthes alata* (Hook.) D. Dietr.

*Prenanthes lessingii* Hultén

*Sonchus hastatus* Less.

***Nestotus* Urbatsch & Neubig [FNA20, HC2]**

Sida. 21: 1650. 2005.

mock goldenweed

**Nestotus stenophyllus** (A. Gray) R.P. Roberts, Urbatsch & Neubig [FNA20, HC2]

Sida. 21: 1652.

narrowleaf goldenweed

*Haplopappus stenophyllus* A. Gray [HC]

*Stenotus stenophyllus* (A. Gray) Greene

**Nothocalais** [FNA19, HC2]

Bull. Calif. Acad. Sci. 2: 54. 1886.

false dandelion

**Nothocalais alpestris** (A. Gray) K.L. Chambers [FNA19, HC2]

Contr. Dudley Herb. 5: 66.

alpine lake agoseris

*Agoseris alpestris* (A. Gray) Greene

*Agoseris barbellulata* Greene

*Microseris alpestris* (A. Gray) Q. Jones ex Cronquist [HC]

**Nothocalais troximoides** (A. Gray) Greene [FNA19, HC2]

Bull. Calif. Acad. Sci. 2: 55.

weevil prairie dandelion, false agoseris

*Microseris troximoides* A. Gray [HC]

*Scorzonella troximoides* (A. Gray) Jeps.

**Onopordum** [FNA19, HC, HC2]

Sp. Pl. 2: 827. 1753; Gen. Pl. ed. 5, 359. 1754.

cotton-thistle

**Onopordum acanthium** L. [FNA19, HC, HC2]

Sp. Pl. 2: 827.

cotton thistle, Scotch thistle, Scots thistle

ssp. **acanthium** [FNA19, HC2]

Scotch thistle, cotton thistle, Scots thistle

**Oreostemma** [FNA20, HC2]

Pittonia. 4: 224. 1900.

mountain-crown

**Oreostemma alpigenum** (Torr. & A. Gray) Greene [FNA20, HC2]

Pittonia. 4: 224.

alpine aster

*Aster alpigenus* (Torr. & A. Gray) A. Gray [HC]

var. **alpigenum** [FNA20, HC2]

Pittonia. 4: 224.

alpine aster, tundra mountaincrown

*Aster alpigenus* (Torr. & A. Gray) A. Gray ssp. *alpigenus*

*Aster alpigenus* (Torr. & A. Gray) A. Gray var. *alpigenus* [HC]

**Packera** [FNA20, HC2]

Bot. Not. 128: 520. 1976.

butterweed, groundsel

**Packera bolanderi** (A. Gray) W.A. Weber & Á. Löve [FNA20, HC2]

Phytologia. 49: 45.

Bolander's groundsel

*Senecio bolanderi* A. Gray [HC]

var. **harfordii** (Greenm.) Trock & T.M. Barkley [FNA20, HC2]

Sida. 18: 386.

Harford's groundsel, Harford's ragwort

*Senecio bolanderi* A. Gray var. *harfordii* (Greenm.) T.M. Barkley [HC]

*Senecio harfordii* Greenm.

**Packera cana** (Hook.) W.A. Weber & Á. Löve [FNA20, HC2]

Phytologia. 49: 46.

woolly groundsel

*Senecio canus* Hook. [HC]

*Senecio howellii* Greene

*Senecio purshianus* Nutt.

**Packera contermina** (Greenm.) J.F. Bain [FNA20, HC2]

Novon. 9: 457.

dwarf arctic butterweed

*Packera cymbalaria* (Pursh) W.A. Weber & Á. Löve [FNA], misapplied

In Washington, <i>Senecio cymbalaria</i>, now <i>Packera cymbalaria</i>, are misapplied names referential to a species distributed in the Arctic.

FNA20: "<i>Packera contermina</i> grows in rocky areas and produces relatively short rhizomes and abundant thin fibrous roots. In mesic sites, the rhizomes are more robust and the fibrous roots are fewer. This taxon has been treated as part of <i>P. cymbalaria</i> or <i>P. subnuda</i>. Morphologic and cytologic data lend support to its recognition at species rank."

**Packera flettii** (Wiegand) W.A. Weber & Á. Löve [FNA20, HC2]

Phytologia. 49: 46.

Flett's groundsel

*Senecio flettii* Wiegand [HC]

**Packera indecora** (Greene) Á. Löve & D. Löve [FNA20, HC2]

Bot. Not. 128: 520.

rayless mountain butterweed, elegant groundsel

*Senecio indecorus* Greene [HC]

**Packera macounii** (Greene) W.A. Weber & Á. Löve [FNA20, HC2]

Phytologia. 49: 47.

Puget butterweed, long-rayed groundsel, Macoun's groundsel, Siskiyou Mountain ragwort

*Senecio macounii* Greene [HC]

FNA20: "Packera macounii is similar in overall morphology to *P. cana*. Leaves of *P. macounii* are narrower and frequently revolute. It is often cited as being collected on serpentine soils; it is not restricted to them. *Senecio fastigiatus* Nuttall (1840) is a later homonym of *S. fastigiatus* Schweinitz ex Elliott (1823), a name of uncertain application." Chambers and Sundberg (2000) note this species is weakly separated from *S. canus* in sw OR

**Packera pauciflora** (Pursh) Á. Löve & D. Löve [FNA20, HC2]

Bot. Not. 128: 520.

rayless alpine butterweed, rayless alpine groundsel

*Senecio pauciflorus* Pursh [HC]

FNA20: "Heads of *Packera pauciflora* are usually discoid. Its range and habitat overlap those of *P. indecora*; the two can be difficult to distinguish."

**Packera paupercula** (Michx.) Á. Löve & D. Löve [FNA20, HC2]

Bot. Not. 128: 520.

Canadian butterweed, balsam groundsel

*Senecio pauperculus* Michx. [HC]

*Senecio pauperulus* Michx. var. *thompsoniensis* (Greenm.) B. Boivin [HC]

FNA20: "Ecologically and morphologically, *Packera pauperula* is the most variable species of the genus in North America. Some "phases" have been treated as separate species, subspecies, varieties, forms, and races. Variation within *P. pauperula* hints at some interesting evolutionary relationships; characteristics used to separate taxa overlap. Much of the morphologic variation in this species may be due to hybridization and introgression. I do not recognize any of the infraspecific taxa that have been proposed."

***Packera porteri* (Greene) C. Jeffrey [FNA20, HC2]**

Kew Bull. 47: 101.

Porter's groundsel

*Senecio porteri* Greene [HC]

FNA20: "Multiple collections of *Packera porteri* are known from Colorado; single collections are known from Oregon (1899; collector indicated few plants were seen) and Washington (1996)." Specimen at Kansas State University.

***Packera pseudaura* (Rydb.) W.A. Weber & Á. Löve [FNA20, HC2]**

Phytologia. 49: 48.

streambank butterweed

*Senecio pseudaura* Rydb. [HC]

**var. *pseudaura* [FNA20, HC2]**

Phytologia. 49: 48.

stream bank butterweed, falsegold groundsel

*Senecio pseudaura* Rydb. ssp. *pseudaura*

*Senecio pseudaura* Rydb. var. *pseudaura* [HC]

***Packera streptanthifolia* (Greene) W.A. Weber & Á. Löve [FNA20, HC2]**

Phytologia. 49: 48.

Rocky Mountain butterweed, cleftleaf groundsel, Rocky Mountain groundsel

*Packera cymbalariaoides* W.A. Weber & A. Löve, invalidly published

*Senecio cymbalariaoides* Nutt. var. *suksdorfii* (Greenm.) M. Peck

*Senecio leonardii* Rydb.

*Senecio streptanthifolius* Greene [HC]

*Senecio streptanthifolius* Greene var. *laetiflorus* (Greene) J.F. Bain

*Senecio streptanthifolius* Greene var. *wallowensis* J.F. Bain

FNA20: "*Packera streptanthifolia* is widespread and variable throughout the Western Cordillera. It includes weakly defined phases that have been treated as distinct species or as varieties. Characteristics used to delimit those taxa often overlap and are difficult to score; some "phases" grade into each other. Northern populations are sometimes segregated as a distinct taxon (e.g., *Senecio streptanthifolia* var. *borealis*; J. F. Bain 1988)." Chambers and Sundberg (2000) follow Bain (1988) in recognizing varieties, but here we do not, following Douglas et al. (1998) and the species concept of Weber

***Packera subnuda* (DC.) Trock & T.M. Barkley [FNA20, HC2]**

Sida. 18: 635.

alpine meadow butterweed, few-leaved groundsel

*Packera bukii* Trock & T.M. Barkley

*Senecio aureus* L. var. *subnudus* (DC.) A. Gray

*Senecio cymbalariaoides* Buek [HC]

*Senecio subnudus* DC.

**var. *subnuda* [FNA20, HC2]**

Sida. 18: 635.

cleftleaf groundsel

*Packera ovina* (Greene) J.F. Bain

*Senecio ovinus* Greene

FNA20: "Plants of *Packera subnuda* var. *subnuda* are scapiform and usually have a single head."

***Petasites*** [FNA20, HC, HC2]

Gard. Dict. Abr. ed. 4. vol. 3. 1754.  
butterbur, coltsfoot

***Petasites frigidus* (L.) Fr.** [FNA20, HC, HC2]

Summa Veg. Scand. 182.  
alpine butterbur, sweet coltsfoot

**var. *frigidus*** [FNA20, HC2]

Summa Veg. Scand. 182.  
alpine butterbur, arctic butterbur, sweet coltsfoot

*Petasites frigidus* (L.) Fr. var. *nivalis* (Greene) Cronquist [HC]

**var. *palmatus* (Aiton) Cronquist** [FNA20, HC, HC2]

Rhodora. 48: 124.  
western coltsfoot

*Nardosmia palmata* (Aiton) Hook.

*Petasites arcticus* A.E. Porsild

*Petasites frigidus* (L.) Fr. ssp. *arcticus* (A.E. Porsild) Cody

*Petasites palmatus* (Aiton) A. Gray

*Petasites palmatus* (Aiton) A. Gray ssp. *speciosus* (Nutt.) Toman

*Petasites speciosus* (Nutt.) Piper

*Tussilago palmata* Aiton

**var. *sagittatus* (Pursh) Cherniawsky & R.J. Bayer** [FNA20, HC2]

Canad. J. Bot. 76: 2070.  
arrowhead sweet coltsfoot, arrowleaf coltsfoot

*Petasites sagittatus* (Pursh) A. Gray [HC]

The taxonomy of this complex, including *Petasites frigidus* and *P. sagittatus*, is disputed. Cherniawsky and Bayer (1998a,b,c) have shown the group has diverged only recently, and proposed a series of varieties with broadly overlapping ranges. We prefer the classification of H&C and JPM, and maintain *P. sagittatus* as a full species.

**var. *xvitifolius* (Greene) Cherniawsky & R.J. Bayer** [FNA20, HC2, KZ99]

Canad. J. Bot. 76: 2072. 1999.  
hybrid coltsfoot, Wenatchee coltsfoot

*Petasites nivalis* Greene ssp. *vitifolius* (Greene) J. Toman

*Petasites trigonophyllus* Greene

*Petasites xvitifolius* Greene [HC]

*Petasites warrenii* H. St. John

FNA20: "Petasites frigidus var. *xvitifolius* often grows in association with one or both putative parents (*P. frigidus* var. *palmatus* and *P. frigidus* var. *sagittatus*)."  
Bogle (1961, 1968) produced this hybrid through artificial crosses.

***Petasites japonicus* (Siebold & Zucc.) Maxim.** [HC2, ILBC1]

Award 34th Demidovian Prize 212.  
Japanese sweet coltsfoot

*Petasites japonicus* (Siebold & Zucc.) Maxim. var. *giganteus* (F. Schmidt ex Trautv.) G. Nicholson [HC2]

*Petasites japonicus* (Siebold & Zucc.) Maxim. var. *japonicus*

***Picris*** [FNA19, HC2]

Sp. Pl. 2: 792. 1753; Gen. Pl. ed. 5, 347. 1754.  
oxtongue, picris

***Picris hieracioides* L.** [FNA19, HC2]

Sp. Pl. 2: 792.  
hawkweed ox-tongue

*Picris hieracioides* L. ssp. *hieracioides*

***Pleiacanthus* [FNA19, HC2]**

Fl. Rocky Mts. 1069. 1917.  
skeletonweed

***Pleiacanthus spinosus* (Nutt.) Rydb. [FNA19, HC2]**

Fl. Rocky Mts. 1069.  
thorny skeletonweed

*Lygodesmia spinosa* Nutt. [HC]

Collected once (2009) in Yakima County. Uncertain as to whether a relictual native stand or an introduction. Closest populations are in central Oregon and southern Idaho.

***Pseudognaphalium* [FNA19, HC2]**

Trudy Bot. Inst. Akad. Nauk S.S.S.R., Ser. 1. Fl. Sist. Vyssh. Rast. 9: 33. 1950.  
cudweed

***Pseudognaphalium californicum* (DC.) Anderb. [FNA19, HC2]**

Opera Bot. 104: 147.  
California cudweed, California everlasting

*Gnaphalium californicum* DC. [HC]

FNA19 does not show this species occurring in WA. Specimen at WTU from Grays Harbor County (1998).

***Pseudognaphalium luteoalbum* (L.) Hilliard & B.L. Burtt [FNA19, HC2]**

Bot. J. Linn. Soc. 82: 206.  
weedy cudweed, red-tip rabbit-tobacco, jersey rabbit tobacco

*Gnaphalium luteo-album* L. [HC], orthographic variant

*Gnaphalium luteoalbum* L.

***Pseudognaphalium macounii* (Greene) Kartesz [FNA19, HC2]**

Synth. N. Amer. Fl. nomencl. innov. 30.  
sticky cudweed, winded cudweed, Macoun's rabbit-tobacco, Macoun's rabbit tobacco

*Gnaphalium macounii* Greene

*Pseudognaphalium viscosum* (Kunth) Anderb. [FNA19], misapplied

FNA20: "Pseudognaphalium macounii is recognized by its stipitate-glandular, proximally glabrescent stems, bicolor and decurrent leaves, relatively large and many-flowered heads, and hyaline, shiny phyllaries." Reported in WA by Creso (1984); Chambers and Sundberg (2000) separate from viscosum but BC flora lumps them. WTU voucher needs check of identity.

***Pseudognaphalium stramineum* (Kunth) Anderb. [FNA19, HC2]**

Opera Bot. 104: 148.  
cotton batting cudweed

*Gnaphalium chilense* Spreng. [HC]

*Gnaphalium stramineum* Kunth

*Pseudognaphalium stramineum* (Kunth) W.A. Weber, invalid name

FNA19 lists this species as occurring in WA. FNA19: "Pseudognaphalium stramineum is probably native from South America to western North America; it is adventive in sandy fields on the Atlantic coastal plain, where it flowers May?Aug."

***Pseudognaphalium thermale* (E.E. Nelson) G.L. Nesom [FNA19, HC2]**

Sida. 21: 781.  
slender cudweed, northwestern rabbit-tobacco

*Gnaphalium canescens* DC. ssp. *thermale* (E.E. Nelson) Stebbins & D.J. Keil

*Gnaphalium microcephalum* Nutt. ssp. *thermale* (E.E. Nelson) G.W. Douglas

*Gnaphalium microcephalum* Nutt. var. *thermale* (E.E. Nelson) Cronquist [HC]

*Pseudognaphalium canescens* (DC.) Anderb. ssp. *thermale* (E.E. Nelson) Kartesz

*Pseudognaphalium microcephalum* (Nutt.) Anderb. [FNA19], misapplied

***Psilocarphus* [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 340. 1840.  
woolly-heads, woolly-marbles

***Psilocarphus brevissimus* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 340.  
dwarf woolly-marbles

**var. *brevissimus* [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 340.  
dwarf woollyheads

*Psilocarphus globiferus* Nutt.

FNA19: "Variety *brevissimus* occupies nearly the full range of the genus (uncommon west of the Cascade Range); some occurrences toward the northeast appear to be recent introductions."

***Psilocarphus elatior* (A. Gray) A. Gray [FNA19, HC, HC2]**

Syn. Fl. N. Amer. ed. 2. 1: 448.  
tall woollyheads

*Psilocarphus oregonus* Nutt. var. *elatior* A. Gray

perhaps best treated as a geographic subspecies of *P. brevissimus* (Chambers and Sundberg 2000). FNA19: "*Psilocarphus elatior* occurs west of the Cascade Range from California to Vancouver Island, British Columbia, and in scattered areas eastward (northwestern Montana, mountains surrounding the border area common to Oregon, Washington, and Idaho). Reports of *P. elatior* from Alberta and Saskatchewan were based on relatively erect forms of *P. brevissimus* var. *brevissimus*. *Psilocarphus elatior* has been of conservation concern in Canada (J. M. Illingworth and G. W. Douglas 1994). Where sympatric, *Psilocarphus elatior* tends to inhabit relatively dry or seasonally flooded sites in more mesic coastal or montane climates and *P. brevissimus* var. *brevissimus* occurs mainly in wetter, seasonally inundated sites in semiarid climates. Some specimens appear to be intermediate; further study may show the two taxa to be better treated as varietally distinct. See also under *P. brevissimus* var. *multiflorus*."

***Psilocarphus oregonus* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 341.  
Oregon woollyheads

FNA19: "*Psilocarphus oregonus* occurs from west-central California through most of Oregon to southeastern Washington, western Idaho, and northern Nevada. Relatively narrow-leaved, montane forms of *P. tenellus* account for reports of *P. oregonus* from the southern Sierra Nevada to Baja California; further study may show these to be intermediates between the two taxa."

***Psilocarphus tenellus* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 341.  
slender woollyheads

*Psilocarphus tenellus* Nutt. var. *tenellus* [HC]

***Pyrrocoma* [FNA20, HC2]**

Fl. Bor.-Amer. 1: 306, plate 107. 1833.  
goldenweed

***Pyrrocoma carthamoides* Hook. [FNA20, HC2]**

Fl. Bor.-Amer. 1: 307, plate 107.  
large-flowered goldenweed

*Haplopappus carthamoides* (Hook.) A. Gray [HC]

**var. *carthamoides* [FNA20, HC2]**

Fl. Bor.-Amer. 1: 307, plate 107.  
Columbia goldenweed, rayless goldenweed

*Haplopappus carthamoides* (Hook.) Gray ssp. *carthamoides*

*Haplopappus carthamoides* (Hook.) A. Gray ssp. *rigidus* (Rydb.) H.M. Hall

*Haplopappus carthamoides* (Hook.) Gray var. *carthamoides* [HC]

FNA20: "Variety *carthamoides* is recognized by its relatively robust stems, large leaves and

involucres, and overlapping, oblong to obovate phyllaries."

**var. *cusickii* (A. Gray) Kartesz & Gandhi [FNA20, HC2]**

Phytologia. 71: 60.

narrowhead goldenweed

*Haplopappus carthamoides* (Hook.) A. Gray ssp. *cusickii* (A. Gray) H.M. Hall

*Haplopappus carthamoides* (Hook.) A. Gray var. *cusickii* A. Gray [HC]

FNA20: "Variety *cusickii* is recognized by its generally smaller size, and campanulate to turbinate involucres with loose, lanceolate phyllaries."

***Pyrrocoma hirta* (A. Gray) Greene [FNA20, HC2]**

Erythea. 2: 69.

hairy goldenweed, sticky goldenweed

*Haplopappus hirtus* A. Gray [HC]

*Haplopappus hirtus* A. Gray var. *hirtus* [HC]

**var. *sonchifolia* (Greene) Kartesz & Gandhi [FNA20, HC2]**

Phytologia. 71: 60.

large sticky goldenweed

*Haplopappus hirtus* A. Gray ssp. *sonchifolius* (Greene) H.M. Hall

*Haplopappus hirtus* A. Gray var. *sonchifolius* (Greene) M. Peck [HC]

FNA20: "Variety *sonchifolia* is recognized by its wider leaves and its preference for moist habitats. More study is needed to determine the status of this taxon."

***Pyrrocoma liatiformis* Greene [FNA20, HC2]**

Leafl. Bot. Observ. Crit. 2: 17.

Palouse goldenweed

*Haplopappus integrifolius* Porter ex A. Gray ssp. *liatiformis* (Greene) H.M. Hall

*Haplopappus integrifolius* Porter ex A. Gray ssp. *scaberulus* (Greene) H.M. Hall

*Haplopappus liatiformis* (Greene) H. St. John [HC]

*Haplopappus racemosus* (Nutt.) Torr. ssp. *liatiformis* (Greene) D.D. Keck

FNA20: "Pyrrocoma liatiformis is one of the dominants of virgin Palouse prairies and appears to be threatened. It is recognized by its hirsute stems, leaves, and phyllaries, and the small, pedunculate heads."

***Pyrrocoma scaberula* Greene [HC2]**

Leaflets of Botanical Observation and Criticism 2(1): 19.

palouse goldenweed

***Rainiera* [FNA20, HC2]**

Pittonia. 3: 291. 1898.

rainiera

***Rainiera stricta* (Greene) Greene [FNA20, HC2]**

Pittonia. 3: 291.

false silverback

*Luina stricta* (Greene) B.L. Rob. [HC]

***Ratibida* [FNA21, HC, HC2]**

Fl. Ludov. 73. 1817.

prairie coneflower, Mexican-hat

***Rhaponticum* [HC2], conserved name**

hardheads, maral root

***Rhaponticum repens* (L.) Hidalgo [HC2]**

Ann. Bot. (Oxford) n.s., 97(5): 714.

hardheads, creeping knapweed, Russian knapweed

*Acroptilon repens* (L.) DC. [FNA19]

*Centaurea repens* L. [HC]

FNA19: "In most American floristic literature *Acroptilon* has been included within *Centaurea*, from which it differs by the subbasal rather than lateral attachment scars on the cypselae and the absence of sterile outer florets. The chromosome base number. = 13 is higher than that in most species of *Centaurea* in the strict sense. Molecular phylogenetic studies of the relationships of Cynareae genera (A. Susanna et al. 1995) support the segregation of *Acroptilon* from *Centaurea*."

***Rigiopappus* [FNA20, HC, HC2]**

Proc. Amer. Acad. Arts. 6: 548. 1865.

***Rigiopappus leptocladus* A. Gray [FNA20, HC, HC2]**

Proc. Amer. Acad. Arts. 6: 548.  
bristlehead, false wireweed

***Rudbeckia* [FNA21, HC, HC2]**

Sp. Pl. 2: 906. 1753; Gen. Pl. ed. 5, 387. 1754.  
coneflower, rudbeckia

***Rudbeckia alpicola* Piper [FNA21, HC2]**

Erythea. 7: 173.  
showy black-eyed Susan, Washington showy black-eyed Susan, Wenatchee Mountain showy black-eyed Susan

*Rudbeckia occidentalis* Nutt. var. *alpicola* (Piper) Cronquist [HC]

***Rudbeckia hirta* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 907.  
black-eyed Susan

**var. *pulcherrima* Farw. [FNA21, HC, HC2]**

Rep. (Annual) Michigan Acad. Sci. 6: 209.  
black-eyed Susan

***Rudbeckia laciniata* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 906.  
green-headed black-eyed Susan, tall black-eyed Susan

**var. *ampla* (A. Nelson) Cronquist [FNA21, HC, HC2]**

Vasc. Pl. Pacif. N.W. 5: 280.  
tall black-eyed Susan

*Rudbeckia ampla* A. Nelson

FNA21: "Cultivars of *Rudbeckia laciniata* are grown as ornamentals. The cultivar "?golden-glow'" is widely planted and occasionally escapes cultivation."

***Rudbeckia occidentalis* Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 355.  
western black-eyed Susan, western chocolate black-eyed Susan  
(see also *Rudbeckia alpicola*)

*Rudbeckia occidentalis* Nutt. var. *occidentalis* [HC]

***Saussurea* [FNA19, HC, HC2]**

Ann. Mus. Natl. Hist. Nat. 16: 156, 196, plates 10?13. 1810.  
saw-wort

***Saussurea americana* D.C. Eaton [FNA19, HC, HC2]**

Bot. Gaz. 6: 283.  
American sawwort

***Senecio* [FNA20, HC, HC2]**

Sp. Pl. 2: 866. 1753; Gen. Pl. ed. 5, 373. 1754.  
butterweed, groundsel, ragwort  
(see also *Jacobaea*, *Packera*)

***Senecio elmeri*** Piper [FNA20, HC, HC2]

Erythea. 7: 173.

Elmer's ragwort

***Senecio fremontii*** Torr. & A. Gray [FNA20, HC, HC2]

Fl. N. Amer. 2: 445.

dwarf mountain butterweed

*Senecio ductoris* Piper

**var. *fremontii*** [FNA20, HC, HC2]

Fl. N. Amer. 2: 445.

dwarf mountain groundsel

***Senecio hydrophilooides*** Rydb. [FNA20, HC2]

Mem. New York Bot. Gard. 1: 441.

sweet marsh butterweed, stout meadow groundsel

*Senecio foetidus* Howell [HC]

*Senecio foetidus* Howell var. *foetidus* [HC]

*Senecio foetidus* Howell var. *hydrophilooides* (Rydb.) T.M. Barkley ex Cronquist [HC]

*Senecio oreganus* Howell

FNA20: "Plants of *Senecio hydrophilooides* from toward the western end of the range tend to have the heads more or less congested and eradiate and stems loosely clustered; plants from toward the eastern edge tend to have heads loosely arrayed and radiate and stems single. The two forms have been recognized as weakly defined species (or varieties), the former as *Senecio foetidus* and the latter as *S. hydrophilooides*. They intergrade so completely that they are best treated as a single, variable taxon. The use of the epithet *foetidus* for the broadly conceived single species was based on a bibliographic misunderstanding; the correct epithet is *hydrophilooides* (T. M. Barkley 1978; A. Cronquist 1994). In 1900, Thomas Howell gave the name *Senecio oreganus* to a collection from Lake Labish, near Salem, Oregon. The area has seen much disturbance and development since Howell's time, and the plant appears to be extinct in the region. The collection is difficult to exclude from *S. hydrophilooides*, and the collection is here regarded as an odd outlier of *S. hydrophilooides*, which is known chiefly from east of the Cascade uplift. Howell's collection and therefore the name *S. oreganus* also have been treated within *S. sphaerocephalus* (T. M. Barkley 1978; A. Cronquist 1955); that attribution appears to be in error. The "type" materials are now in the herbarium of Oregon State University in Corvallis."

***Senecio hydrophilus*** Nutt. [FNA20, HC, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 411.

alkali marsh butterweed, water groundsel, alkali marsh ragwort

*Senecio hydrophilus* Nutt. var. *pacifica* Greene

***Senecio integerrimus*** Nutt. [FNA20, HC, HC2]

Gen. N. Amer. Pl. 2: 165.

one-stemmed butterweed, western groundsel

**var. *exaltatus*** (Nutt.) Cronquist [FNA20, HC, HC2]

Leafl. W. Bot. 6: 48.

one-stemmed butterweed, lambs-tongue groundsel, tall western groundsel

*Senecio integerrimus* Nutt. var. *vaseyi* (Greenm.) Cronquist [HC]

*Senecio vaseyi* Greenm.

FNA20: "Variety *< i>exaltatus</i>* is the most widespread and variable variety of the species. Eradiate plants of var. *< i>exaltatus</i>* have been recognized as var. *< i>vaseyi</i>*; there appears to be no populational integrity to the eradiate condition." Chambers and Sundberg (2000) note this is very similar to *< i>S. hydrophilooides</i>* and differs only in the pubescence.

**var. *ochroleucus*** (A. Gray) Cronquist [FNA20, HC, HC2]

Leafl. W. Bot. 6: 48.

white western groundsel

*Senecio exaltatus* Nutt. ssp. *ochraceus* Piper

Chambers and Sundberg (2000) note the flower color cannot be determined on older herbarium

sheets, but the cordate or sub-cordate leaves are unique

***Senecio lugens* Richardson [FNA20, HC, HC2]**

Narr. Journey Polar Sea. 748.  
black-tipped groundsel

*Senecio integrerrimus* Nutt. var. *lugens* (Richardson) B. Boivin

***Senecio neowebsteri* S.F. Blake [FNA20, HC, HC2]**

Leafl. W. Bot. 8: 143.  
Olympic Mountain ragwort

***Senecio serra* Hook. [FNA20, HC, HC2]**

Fl. Bor.-Amer. 1: 333.  
tall butterweed, butterweed groundsel

**var. *serra* [FNA20, HC2]**

Fl. Bor.-Amer. 1: 333.  
tall butterweed, butterweed groundsel

*Senecio andinus* Nutt.

*Senecio lanceolatus* Torr. & A. Gray

*Senecio millikenii* Eastw.

*Senecio serra* Hook. var. *altior* Jeps.

*Senecio solidago* Rydb.

***Senecio sylvaticus* L. [FNA20, HC, HC2]**

Sp. Pl. 2: 868.  
wood groundsel, woodland groundsel, woodland ragwort

FNA20: "Senecio sylvaticus is a Eurasian weed that favors cool, wet climates. It is well established in coastal areas of the Pacific Coast and in parts of Newfoundland and Quebec; elsewhere in the flora, it appears to be sporadic."

***Senecio triangularis* Hook. [FNA20, HC, HC2]**

Fl. Bor.-Amer. 1: 332, plate 115.  
arrowleaf groundsel, arrowleaf ragwort

*Senecio triangularis* Hook. var. *angustifolius* G.N. Jones [HC]

FNA20: "Plants of *Senecio triangularis* with narrow, subentire leaves that taper to the petioles are occasionally encountered in acid bogs in Oregon and Washington and less frequently elsewhere. They are regarded as edaphic variants; they have been recognized as var. *angustifolius*." Var. *angustifolius* of southern OR & CA is distinct (Chambers and Sundberg, 2000).

**var. *triangularis* [HC, HC2]**

arrowleaf groundsel, arrowleaf ragwort

***Senecio viscosus* L. [FNA20, HC2]**

Sp. Pl. 2: 868.  
sticky ragwort

FNA 20: "Senecio viscosus is a smelly, Eurasian weed now widely scattered in areas of cool damp climates, often as a casual waif. The viscid hairs trap wind-blown particles of sand, dust, and soot, which give the surfaces varying textures and colors."

***Senecio vulgaris* L. [FNA20, HC, HC2]**

Sp. Pl. 2: 867.  
common groundsel, old man in the spring

***Sericocarpus* [FNA20, HC2]**

Gen. Sp. Aster. 10, 148. 1832.  
white-topped aster

***Sericocarpus oregonensis* Nutt. [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 302.  
Oregon white-topped aster

*Aster oregonensis* (Nutt.) Cronquist [HC]

ssp. *oregonensis* [FNA20, HC2]

Trans. Amer. Philos. Soc., n. s. 7: 302.

Oregon white-topped aster

*Aster oregonensis* (Nutt.) Cronquist ssp. *oregonensis*

*Sericocarpus oregonensis* Nutt. var. *oregonensis*

***Sericocarpus rigidus* Lindl. [FNA20, HC2]**

Fl. Bor. Amer. 2: 14.

Columbian white-topped aster, Columbian whitetop aster, rigid white-topped aster

*Aster curtus* Cronquist [HC]

***Silybum* [FNA19, HC, HC2]**

Fam. Pl. 2: 116, 605. 1763.

milk-thistle

***Silybum marianum* (L.) Gaertn. [FNA19, HC, HC2]**

Fruct. Sem. Pl. 2: 378.

milk-thistle

***Solidago* [FNA20, HC, HC2]**

Sp. Pl. 2: 878. 1753; Gen. Pl. ed. 5, 374. 1754.

goldenrod

(see also *Euthamia*)

***Solidago altissima* L. [FNA20, HC2]**

Sp. Pl. 2: 878.

Great Plains goldenrod, late goldenrod

ssp. *altissima* [FNA20]

late goldenrod

Known from several locations in Grant County.

ssp. *gilvacanescens* (Rydb.) Semple [FNA20, HC2]

Sida. 20: 1606.

Great Plains goldenrod, tall goldenrod

*Solidago canadensis* L. var. *gilvacanescens* Rydb. [HC]

***Solidago elongata* Nutt. [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 327.

Cascade Canada goldenrod, West Coast goldenrod

*Solidago canadensis* L. ssp. *elongata* (Nutt.) D.D. Keck

*Solidago lepida* DC. var. *caurina* (Piper) M. Peck

*Solidago lepida* DC. var. *elongata* (Nutt.) Fernald

FNA20: "This species can be similar to *S. lepida*, which usually has much larger distal cauline leaves."

***Solidago lepida* DC. [FNA20, HC2]**

Prod. 5: 339.

western Canada goldenrod

*Solidago canadensis* L. var. *lempira* (DC.) Cronquist

*Solidago canadensis* L. var. *subserrata* (DC.) Cronquist [HC]

FNA20: "Solidago lepida is the only member of subsect. *Triplinerviae* that is nearly always minutely, sometimes sparsely, stipitate-glandular in the arrays. These glands can be very small and visible only with 30?70x magnification. The enlarged head of the stalked gland may be little bigger than a pollen grain. The species occurs in the west from the Aleutian Islands and central Alaska south just into northern California, and in the Rocky Mountains to Arizona and New Mexico. Its range extends across Canada to Newfoundland, the Gaspé (Quebec), and northern New Brunswick. It also appears to be introduced at scattered locations across the Canadian prairies."

**var. *leptophylla*** [FNA20, HC2]

In A. P. de Candolle and A. L. P. P. de Candolle, Prodr. 5: 339.  
western Canada goldenrod

FNA20: "Variety leptophylla can be difficult to distinguish from *Solidago elongata* in the Cascades and coastal areas of southern British Columbia and Washington. Involucre height increases with ploidy level."

**var. *salebrosa* (Piper) Semple** [FNA20, HC2]

Sida. 20: 1611.  
Canada goldenrod, meadow goldenrod, Rocky Mountains Canada goldenrod

*Solidago canadensis* L. ssp. *salebrosa* (Piper) D.D. Keck

*Solidago canadensis* L. var. *salebrosa* (Piper) M.E. Jones [HC]

*Solidago gigantea* Aiton [FNA20, HC, HC2], misapplied

FNA20: "FNA20: "Variety *salebrosa* strongly resembles *Solidago canadensis*, and is found throughout most of the Rocky Mountains in the United States and adjacent Canada. It has been included in *S. canadensis* by many authors (e.g., A. Cronquist 1994). In extreme forms the array is broader than tall with long, arching proximal branches. Hairier plants can be similar in appearance to *S. altissima*; the latter is usually not glandular and is much hairier. Glabrate plants of var. *salebrosa* can be difficult to distinguish from hexaploid *S. gigantea* near and in the mountains from Alberta south to New Mexico. Glabrate plants in the mountains often treated as *S. gigantea* are glandular and belong in *S. leptocephala* var. *salebrosa*. Small-headed diploids found in the Rocky Mountains from southern British Columbia to Colorado are usually sparsely glandular and could be confused with short-array forms of *S. elongata*." Check WTU colls for specimens of *S. altissima*, reported N to BC by Semple (1993)

***Solidago missouriensis* Nutt.** [FNA20, HC, HC2]

J. Acad. Nat. Sci. Philadelphia. 7: 32.  
Missouri goldenrod

*Solidago missouriensis* Nutt. var. *extraria* A. Gray [HC]

*Solidago missouriensis* Nutt. var. *fasciculata* Holz. [HC]

*Solidago missouriensis* Nutt. var. *missouriensis* [HC]

*Solidago missouriensis* Nutt. var. *tolmieana* (A. Gray) Cronquist [HC]

FNA20: "*Solidago missouriensis* was often introduced along railroad lines farther east. It is a highly variable species. In the east, it can be similar to *S. juncea* and is not always easily distinguished where ranges overlap. In the west, it can similar to smaller plants of *S. spectabilis*. It is distinguished from the related species by its usually 3-nerved proximal leaves and the usually thin, elongate rhizomes. Across the prairies the species is known to be diploid only ( $2n = 18$ ). In the Rocky Mountains, tetraploids ( $2n = 36$ ) are common, the diploids infrequent. A number of varieties have been described. Shorter, often larger-headed plants (tetraploids when known) from the Rocky Mountains have been treated as var. *missouriensis* (including var. *extraria*). Taller, more leafy-stemmed plants, mostly from the eastern half of the range, but occasionally west to Washington, have been treated as var. *fasciculata*. Plants from Arizona, Colorado, and New Mexico with long, linear leaves have been treated as var. *tenuissima*. Larger-headed plants with narrow bracts from prairies west of the Cascades in Oregon and Washington have been treated as var. *tolmieana*. A. Cronquist (1994) opted not to recognize varieties, noting that all appeared to grade continuously into each other. A detailed study of the species is needed."

***Solidago multiradiata* Aiton** [FNA20, HC, HC2]

Hort. Kew. 3: 218.  
northern goldenrod, Rocky Mountain goldenrod

*Solidago multiradiata* Aiton ssp. *scopulorum* (A. Gray) W.A. Weber

*Solidago multiradiata* Aiton var. *scopulorum* A. Gray [HC]

FNA20: "*Solidago multiradiata* is the North American species most closely related to *S. virgaurea*, the type species of the genus, native to mostly arctic and alpine regions of Eurasia. Plants of *S. multiradiata* from the Rocky Mountains have been treated as var. *scopulorum*; they differ so little from those of other parts of the range that recognition of the variety without further support does not appear justified."

***Solidago simplex* Kunth** [FNA20, HC2]

Nov. Gen. Sp. 4(fol.): 81. 1818; 4(qto.): 103.

sticky goldenrod

var. *nana* (A. Gray) G.S. Ringius [FNA20, HC2]

Phytologia. 70: 397.

dwarf goldenrod

*Solidago spathulata* DC. var. *nana* (A. Gray) Cronquist [HC]

var. *simplex* [FNA20, HC2]

In A. von Humboldt et al., Nov. Gen. Sp. 4(fol.): 81. 1818; 4(qto.): 103.

sticky goldenrod

*Solidago decumbens* Greene

*Solidago spathulata* DC. ssp. *glutinosa* (Nutt.) D.D. Keck

*Solidago spathulata* DC. var. *neomexicana* (A. Gray) Cronquist [HC]

FNA20: "Variety simplex is found in western North America and is disjunct along the shores of the upper Great Lakes and in southern Quebec."

***Solidago spathulata* DC. [FNA20, HC, HC2]**

Prod. 5: 339.

coast goldenrod, dune goldenrod

*Solidago spathulata* DC. var. *spathulata* [HC]

***Soliva* [FNA19, HC2]**

Fl. Peruv. Prod. 113, plate 24. 1794.

burrweed

***Soliva sessilis* Ruiz & Pav. [FNA19, HC2]**

Syst. Veg. Fl. Peruv. Chil. 113, plate 24.

lawn burrweed, common soliva, prickly soliva

*Soliva pterosperma* (Juss.) Less.

***Sonchus* [FNA19, HC, HC2]**

Sp. Pl. 2: 793. 1753; Gen. Pl. ed. 5, 347. 1754.

sow-thistle

***Sonchus arvensis* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 793.

field sow-thistle, perennial sow-thistle

ssp. *arvensis* [FNA19, HC2]

corn sow-thistle, field sow-thistle, perennial sow-thistle

*Sonchus arvensis* L. var. *arvensis*

ssp. *uliginosus* (M. Bieb.) Nyman [FNA19, HC2]

Consp. Fl. Eur. 433.

field sow-thistle, marsh sow-thistle, wet ground sow-thistle

*Sonchus arvensis* L. var. *glabrescens* Günther

*Sonchus uliginosus* M. Bieb. [HC]

***Sonchus asper* (L.) Hill [FNA19, HC, HC2]**

Herb. Brit. 1: 47.

prickly sow-thistle, spiny leaf sow-thistle

*Sonchus oleraceus* L. var. *asper* L.

ssp. *asper* [HC2]

***Sonchus oleraceus* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 794.

common sow-thistle

***Stenotus* [FNA20, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 334. 1840.  
mock goldenweed

***Stenotus lanuginosus* (A. Gray) Greene [FNA20, HC2]**

Erythea. 2: 72.

woolly goldenweed

*Haplopappus lanuginosus* A. Gray [HC]

**var. *lanuginosus* [FNA20, HC2]**

Erythea. 2: 72.

woolly goldenweed, woolly stenotus

*Haplopappus lanuginosus* Gray var. *lanuginosus* [HC]

***Stephanomeria* [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 427. 1841.  
wirelettuce

***Stephanomeria exigua* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 428.  
small wirelettuce

**ssp. *exigua* [FNA19]**

Trans. Amer. Philos. Soc., n. s. 7: 428.  
skeletonplant, small wirelettuce

***Stephanomeria paniculata* Nutt. [FNA19, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 428.  
stiff-branched stephanomeria, stiff-branched wirelettuce

***Stephanomeria tenuifolia* (Raf.) H.M. Hall [FNA19, HC, HC2]**

Univ. Calif. Publ. Bot. 3: 256.  
wire lettuce, narrowleaf stephanomeria

*Stephanomeria minor* (Hook.) Nutt. var. *minor*

*Stephanomeria tenuifolia* (Raf.) H.M. Hall var. *myrioclada* (D.C. Eaton) Cronquist [HC]

*Stephanomeria tenuifolia* (Raf.) H.M. Hall var. *tenuifolia* [HC]

FNA19: "Stephanomeria tenuifolia is distributed over an immense region and is the most widespread species of the genus. It shows remarkable variability in the form and dimensions of its stems and branches. Plants described as *S. myrioclada*, from the northeasternmost corner of Nevada, present an architecture of relatively numerous, almost threadlike, densely crowded stems (1.5?4 dm) and branches with an irregularly dichotomous pattern. Continuous variation occurs from this form to another in the same region and elsewhere in which the stems are longer (3?7 dm), sparingly branched, and flexuous. The extreme variability in vegetative architecture may be adaptive and deserves further study."

***Symphyotrichum* [FNA20, HC2]**

Gen. Sp. Aster. 9, 135. 1832.  
aster

***Symphyotrichum xamethystinum* (Nutt.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 294. 1995.  
amethyst aster, hybrid aster

*Aster amethystinus* Nutt.

FNA20: "Symphyotrichum xamethystinum is the F 1 hybrid between *S. ericoides* and *S. novae-angliae*, encountered sometimes throughout the area where the two parental species co-occur. It is morphologically intermediate; it has non-spiny, sparsely stipitate-glandular phyllaries and rose-violet rays in mid-sized heads. Forma leucerythros Bemis and forma leucos Bemis have been described within this hybrid and may represent recombinants or normal population color variants."

***Symphyotrichum ascendens* (Lindl.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 276.  
intermountain aster, long-leaved aster, western American aster, western aster

*Aster adscendens* Lindl.

*Aster ascendens* Lindl.

*Aster chilensis* Nees ssp. *adscendens* (Lindl.) Cronquist [HC]

*Aster adscendens* orth. error in Abrams FNA20: "Symphyotrichum ascendens is widely distributed in the Great Basin. It is an allopolyploid derived from the hybrid between *S. spathulatum* ( $x = 8$ ) and *S. falcatum* ( $x = 5$ ). Chromosome numbers differ markedly in their geographic distribution,  $2n = 26$  prevailing in the southwestern part of the range, and  $2n = 52$  in the northeastern part (G. A. Allen 1985) Backcrosses to both parental species or hybrids with related taxa are sometimes seen where the ranges overlap."

***Symphyotrichum boreale* (Torr. & A. Gray) Å. Löve & D. Löve [FNA20, HC2]**

Taxon. 31: 358. 1982.

northern bog aster, rush aster, slender white aster

*Aster borealis* (Torr. & A. Gray) Provancher

*Aster junciformis* Rydb. [HC]

*Aster laxiflorus* Lindl. var. *borealis* Torr. & A. Gray

***Symphyotrichum bracteolatum* (Nutt.) G.L. Nesom [HC2, JPM2]**

Phytologia 77(3): 276.

bracted aster, Eaton's aster, Oregon aster

*Aster eatonii* (A. Gray) Howell [HC]

*Symphyotrichum eatonii* (A. Gray) G.L. Nesom [FNA20]

Jepson 2nd: "The name *S. bracteolatum* has nomenclatural priority over *S. eatonii* (Brummitt 2011 Taxon 60:230)."

***Symphyotrichum campestre* (Nutt.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 276.

western meadow aster

*Aster campestris* Nutt. [HC]

*Aster campestris* Nutt. var. *bloomeri* (A. Gray) A. Gray [HC]

*Aster campestris* Nutt. var. *campestris* [HC]

*Symphyotrichum campestre* (Nutt.) G.L. Nesom var. *bloomeri* (A. Gray) G.L. Nesom

*Symphyotrichum campestre* (Nutt.) G.L. Nesom var. *campestris*

FNA20: "Two poorly defined varieties of *Symphyotrichum campestre* have been described. Variety *campestre* has glabrous or sparsely strigose leaves and occurs in southern British Columbia, Colorado, Idaho, Montana, Nevada, Oregon, Washington, and Wyoming. Bloomer? Aster, var. *bloomeri*, has moderately strigose leaves and occurs in California, Nevada, and Oregon. The varieties are not sufficiently distinct to warrant recognition. *Symphyotrichum ×columbianum* (Piper) G. L. Nesom (syn. *Aster columbianus* Piper, *A. multiflorus* Aiton var. *columbianus* (Piper) S. F. Blake, *Virgulus ×columbianus* (Piper) Reveal & Keener) is the hybrid between *S. campestre* and *S. ericoides* subsp. *pansum*."

***Symphyotrichum chilense* (Nees) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 277.

common California aster, Pacific aster

(see also *Symphyotrichum ascendens*, *Symphyotrichum hallii*)

*Aster chilensis* Nees [HC]

*Aster chilensis* Nees ssp. *chilensis* [HC]

*Aster chilensis* Nees var. *chilensis*

*Symphyotrichum chilense* (Nees) G.L. Nesom var. *chilense*

FNA20: "*Symphyotrichum chilense* is restricted to coastal habitats from southwestern British Columbia to central California. It is almost entirely coastal in Oregon, Washington, and southern British Columbia, where it is mainly hexaploid ( $2n = 48$ ). In Oregon, where it is sympatric with *S. subspicatum*, the latter is mainly duodecaploid ( $2n = 96$ ). The distinction does not hold in British Columbia, however, where *S. subspicatum* is both  $2n = 48$  and  $96$ , and where *S. chilense* is less common (G. A. Allen 1984). The species was erroneously thought by Nees to occur in Chile. The plants named *Aster chilensis* var. *medius* Jepson are hybrids of *S. chilense* and *S. lenticum*."

***Symphyotrichum ciliatum* (Ledeb.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 277.

alkali American aster, rayless alkali aster, rayless annual aster

*Aster brachyactis* S.F. Blake [HC]

***Symphyotrichum ×columbianum* (Piper) G.L. Nesom [FNA20, HC2]**

hybrid aster

*Aster columbianus* Piper

***Symphyotrichum ericoides* (L.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 280.

heath-leaved aster, tufted white prairie aster

**var. *pansum* (S.F. Blake) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 280.

little gray aster, tufted white prairie aster, white heath aster

*Aster ericoides* L. ssp. *pansus* (S.F. Blake) A.G. Jones

*Aster ericoides* L. var. *pansus* (S.F. Blake) B. Boivin

*Aster pansus* (S.F. Blake) Cronquist [HC]

FNA21: "Variety *pansum* is reported to be rare in extreme western Kansas and Ontario (where it is introduced), extreme northeastern Arizona, and northwestern New Mexico. It has been introduced along railroads farther east. A. G. Jones (1978) treated this taxon as a subspecies with two varieties. Plants forming clumps with many, erect to arching, stout, usually densely hispido-strigose stems were called var. *pansum*; these occur mostly in British Columbia, Idaho, Oregon, and Washington. Plants in clusters with few, decumbent or ascending, slender, usually sparsely strigose stems were recognized by Jones as var. *stricticaule*; these are encountered mostly on the prairies and in the foothills of the Rocky Mountains from Alberta to Manitoba, Utah, and Wyoming."

***Symphyotrichum foliaceum* (Lindl. ex DC.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 282.

alpine leafybract aster, Canby's leafybract aster, Cusick's American aster, Cusick's aster, Henderson's aster, Kootenai aster, leafy aster, leafy-bracted aster, Parry's aster

*Aster cusickii* A. Gray

*Aster foliaceus* Lindl. ex DC. [HC]

*Aster foliaceus* Lindl. ex DC. var. *apricus* A. Gray [HC]

*Aster foliaceus* Lindl. ex DC. var. *canbyi* A. Gray [HC]

*Aster foliaceus* Lindl. var. *cusickii* (A. Gray) Cronquist [HC]

*Aster foliaceus* Lindl. ex DC. var. *foliaceus* [HC]

*Aster foliaceus* Lindl. ex DC. var. *frondeus* A. Gray

*Aster foliaceus* Lindl. ex DC. var. *lyallii* (A. Gray) Cronquist [HC]

*Aster foliaceus* Lindl. ex DC. var. *parryi* (D.C. Eaton) A. Gray [HC]

*Aster hendersonii* Fernald

*Symphyotrichum cusickii* (A. Gray) G.L. Nesom [FNA20]

*Symphyotrichum foliaceum* (Lindl. ex DC.) G.L. Nesom var. *apricum* (A. Gray) G.L. Nesom [FNA20]

*Symphyotrichum foliaceum* (Lindl. ex DC.) G.L. Nesom var. *canbyi* (A. Gray) G.L. Nesom [FNA20]

*Symphyotrichum foliaceum* (Lindl. ex DC.) G.L. Nesom var. *foliaceum* [FNA20]

*Symphyotrichum foliaceum* (Lindl. ex DC.) G.L. Nesom var. *parryi* (D.C. Eaton) G.L. Nesom [FNA20]

*Symphyotrichum hendersonii* (Fernald) G.L. Nesom [FNA20]

***Symphyotrichum frondosum* (Nutt.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 282.

alkali aster, short rayed aster, short-rayed alkali aster

*Aster frondosus* (Nutt.) Torr. & A. Gray [HC]

*Brachyactis frondosa* (Nutt.) A. Gray

***Symphyotrichum hallii* (A. Gray) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 283.

Hall's aster

*Aster chilensis* Nees ssp. *hallii* (A. Gray) Cronquist [HC]

*Aster hallii* A. Gray

FNA20: "Symphyotrichum hallii is restricted to open habitats of the Puget Trough of western Washington and the Willamette Valley of western Oregon, with outlying stations in the Columbia Gorge and central Washington. Some of the polyploid races appear to be allopoloids involving the sympatric *S. subspicatum*, with larger leaves and fewer, larger heads with violet rays."

***Symphyotrichum jessicae* (Piper) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 283.

Jessica's aster, Palouse aster

*Aster jessicae* Piper [HC]

FNA20: "Symphyotrichum jessicae is known only from the Palouse and Clearwater river drainages of eastern Washington and adjacent northwestern Idaho."

***Symphyotrichum laeve* (L.) Á. Löve & D. Löve [FNA20, HC2]**

Taxon. 31: 359.

Geyer's aster, smooth aster

*Aster laevis* L. [HC]

**var. *geyeri* (A. Gray) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 284.

Geyer's smooth aster

*Aster geyeri* (A. Gray) Howell

*Aster laevis* L. var. *geyeri* A. Gray [HC]

***Symphyotrichum lanceolatum* (Willd.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 284.

marsh aster, western willow aster

**var. *hesperium* (A. Gray) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 284.

lance-leaved aster, panicled aster, white panicled aster

*Aster hesperius* A. Gray [HC]

FNA20: "This variety has been treated mostly as a distinct species in floras. Character ranges overlap considerably with var. *lanceolatum*, and it is often difficult to distinguish the two entities where their distributions overlap. In areas of sympatry, the two taxa hybridize to form septaploid plants ( $2n = 56$ )."

***Symphyotrichum novae-angliae* (L.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 287.

New England aster

*Aster novae-angliae* L. [HC]

FNA20: "Symphyotrichum novae-angliae is escaped from cultivation and introduced in Montana, Oregon, Utah, Washington, and Wyoming, and has been reported as an ephemeral escape in British Columbia. It possibly escaped from cultivation elsewhere. The Michaelmas daisy is widely sold in the horticultural trade, where cultivars have been developed. Forms have been described that correspond to color genetic variants within natural populations (*Aster novae-angliae* forma *roseus* (Desfontaines) Britton; *A. novae-angliae* forma *geneseensis* House); they are not recognized here. *Symphyotrichum novae-angliae* resembles *Canadanthus modestus*, but the ranges of the two do not overlap, and the latter has sparsely hairy cypselae with dark ribs. *Symphyotrichum novae-angliae* hybridizes with *S. ericoides*, forming the F 1 intersectional hybrid *S. ×amethystinum*."

***Symphyotrichum novi-belgii* (L.) G.L. Nesom [FNA20]**

Phytologia. 77: 287.

New York aster

See "A taxonomic revision of the genera *Antennaria* and *Symphyotrichum* (Asteraceae) in British Columbia, Canada, with additional perspectives on the role of taxonomy within the biological sciences" by Jamie Fenneman (<https://open.library.ubc.ca/soa/clrcle/collections/ubctheses/24/items/1.0378325>) for additional information on the occurrence of this species in the Pacific Northwest.

***Symphyotrichum pilosum* (Willd.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 289.

hairy aster

*Aster pilosus* Willd. [HC]

var. *pilosum* [FNA20, HC2]

hairy aster

*Aster pilosus* Willd. var. *pilosus* [HC]

***Sympyotrichum spathulatum* (Lindl.) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 291.

western aster, western mountain aster

*Aster occidentalis* (Nutt.) Torr. & A. Gray [HC]

*Aster occidentalis* (Nutt.) Torr. & A. Gray var. *intermedius* A. Gray [HC]

*Aster occidentalis* (Nutt.) Torr. & A. Gray var. *occidentalis* [HC]

*Aster spathulatus* Lindl. var. *spathulatus*

*Aster vallicola* Greene

*Sympyotrichum spathulatum* (Lindl.) G.L. Nesom var. *intermedium* (A. Gray) G.L. Nesom [FNA20]

*Sympyotrichum spathulatum* (Lindl.) G.L. Nesom var. *spathulatum* [FNA20]

*Sympyotrichum spathulatum* (Lindl.) G.L. Nesom var. *yosemitanum* (A. Gray) G.L. Nesom [FNA20]

***Sympyotrichum subspicatum* (Nees) G.L. Nesom [FNA20, HC2]**

Phytologia. 77: 293.

Douglas' aster, Douglas's aster

*Aster bulteri* Rydb.

*Aster douglasii* Lindl.

*Aster maccallae* Rydb.

*Aster subspicatus* Nees [HC]

*Aster subspicatus* Nees var. *grayi* (Suksd.) Cronquist

*Aster subspicatus* Nees var. *subspicatus*

*Sympyotrichum subspicatum* (Nees) G.L. Nesom var. *grayi* (Suksd.) G.L. Nesom

*Sympyotrichum subspicatum* (Nees) G.L. Nesom var. *subspicatum*

See FNA Volume 20 for description of taxonomic boundaries for this species. FNA20: "Sympyotrichum subspicatum is a weedy, highly polyploid species, probably of allopolyploid derivation from different combinations of species including *S. chilense*, *S. eatonii*, *S. foliaceum*, *S. laeve*, and *S. spathulatum*. Hybrids with *S. hallii* are known from western Oregon. The species passes into *S. foliaceum* in southeastern Alaska."

***Tanacetum* [FNA19, HC, HC2]**

Sp. Pl. 2: 843. 1753; Gen. Pl. ed. 5, 366. 1754.

tansy

(see also *Artemisia*)

***Tanacetum balsamita* L. [FNA19, HC2]**

Sp. Pl. 2: 845.

costmary

*Balsamita major* Desf.

*Chrysanthemum balsamita* (L.) Baillon [HC]

***Tanacetum bipinnatum* (L.) Sch. Bip. [FNA19, HC2]**

Tanaceteen. 48.

camphor tansy, dune tansy

*Tanacetum bipinnatum* (L.) Sch. Bip. ssp. *huronense* (Nutt.) Breitung

*Tanacetum camphoratum* Less.

*Tanacetum douglasii* DC. [HC]

FNA19: "The circumscription of *Tanacetum bipinnatum* adopted here includes not only *T. huronense* (see E. Hultén 1941?1950, vol. 10, 1968) but *T. camphoratum* and *T. douglasii* as well (see D. W. Kyhos and P. H. Raven 1982; C. J. Mickelson and H. H. Iltis 1966). Subspecies *bipinnatum* has been distinguished from ssp. *huronense* by having heads borne singly or 2?4 together versus (1?)3?12(?)20+ in corymbiform arrays, phyllary margins dark brown versus pale brown, and laminae of ray corollas mostly 3?7 mm versus

1?3 mm. Relatively low plants, 10?20(?40 cm) from dune habitats along the southern shore of Lake Athabasca, Saskatchewan, with mostly 1?4, lanate cauline leaves and 1(?) heads per flowering stem have been called *T. huronense* var. *floccosum*."

*Tanacetum parthenium* (L.) Sch. Bip. [FNA19, HC2]

Tanaceen. 55.

featherfew, feverfew

*Chrysanthemum parthenium* (L.) Bernh. [HC]

FNA19:"*Tanacetum parthenium* is widely cultivated throughout North America."

*Tanacetum vulgare* L. [FNA19, HC, HC2]

Sp. Pl. 2: 844.

common tansy

FNA20: "*Tanacetum vulgare* escapes from and/or persists after cultivation. In the flora area, it is naturalized mostly in the northeastern and Pacific Coast states and provinces and sporadically elsewhere."

*Taraxacum* [FNA19, HC, HC2]

Prim. Fl. Holsat. 56. 1780.

dandelion

*Taraxacum alaskanum* Rydb. [FNA19, HC2]

Bull. Torrey Bot. Club. 28: 512.

dwarf alpine dandelion

*Taraxacum kamtschaticum* Dahlst.

First collected in WA in Skagit County in 2019.

*Taraxacum ceratophorum* (Ledeb.) DC. [FNA19, HC, HC2]

Prodri. 7: 146.

horned dandelion

*Taraxacum eriophorum* Rydb. [HC]

*Taraxacum officinale* F.H. Wigg. ssp. *ceratophorum* (Ledeb.) Schinz ex Thell.

*Taraxacum paucisquamsum* M. Peck

*Taraxacum sibiricum* Dahlst.

FNA19: "*Taraxacum ceratophorum* is the most widespread native dandelion in North America, ranging from the low Arctic and boreal zone to the western Cordilleras, in the montane and alpine zones. This complex has been subdivided into many microspecies in North America, most of which appear unworthy of recognition." See Douglas et al. (1998) for review of taxonomic treatments, complicated by apomixis, polyploidy, and hybridization

*Taraxacum erythrospermum* Andrz. ex Besser [FNA19, HC2]

Enum. Pl. 75.

red-seeded dandelion

*Taraxacum laevigatum* (Willd.) DC. [HC]

*Taraxacum laevigatum* (Willd.) DC. var. *erythrospermum* (Andrz. ex Besser) J. Weiss

*Taraxacum officinale* F.H. Wigg. var. *erythrospermum* (Andrz. ex Besser) Bab.

*Taraxacum scanicum* Dahlst.

FNA19: "Early leaves of *Taraxacum erythrospermum* sometimes may be broadly winged along the midvein, making distinction from *T. officinale* difficult; usually, its later leaves become more deeply lobed with time. The name *Taraxacum laevigatum* has been used for *L. erythrospermum* in North America, following H. Handel-Mazzetti (1907). L. H. Shinners (1949) questioned that usage. The name is listed in the index of *Flora Europaea* (A. J. Richards and P. D. Sell 1973) as an unassigned synonym; it could be related to three different entities of sect. Spectabilia. And, it is not mentioned by other modern students of the group. Therefore, (1) given that the North American entity has not been identified with a particular Eurasian taxon; (2) to avoid using a microspecies name such as *T. scanicum*; and (3) despite the lack of typification of the name, I am using *T. erythrospermum* as a place holder until nomenclatural issues are resolved. This clearly associates the taxon with the section to which it belongs."

*Taraxacum officinale* F.H. Wigg. [FNA19, HC, HC2]

Prim. Fl. Holsat. 56.  
common dandelion

*Taraxacum officinale* F.H. Wigg. ssp. *vulgare* (Lam.) Schinz & R. Keller

FNA19: " Taraxacum officinale is the most widespread dandelion in temperate North America, though its abundance decreases in the arid south. It is a familiar weed of lawns and roadsides. It is also the species most commonly used for medicinal and culinary purposes (e.g., E. Small and P. M. Catling 1999).

Phenotypic and genotypic variation of this species have been studied in North America (L. M. King 1993; King and B. A. Schaal 1990; J. C. Lyman and N. C. Ellstrand 1998; O. T. Solbrig 1971; R. J. Taylor 1987), but results of those studies did not lead to the recognition of microspecies.

Specimens of *Taraxacum officinale* with deeply lobed leaves are sometimes difficult to distinguish from those of *T. erythrospermum* when fruits are missing (see also R. J. Taylor 1987). Usually, however, early leaves of the former are much less deeply lobed than those of the latter, which are more consistently lacerate throughout development, though broadly winged initially. The two taxa are easily distinguished in fruit, the red cypselae of *T. erythrospermum* standing out from the dull olive ones of *T. officinale*.

In northeastern North America, *Taraxacum officinale* and *T. lapponicum* often are confused, which has led to reports of the common dandelion farther north than I have been able to verify (it has yet to be collected from the Nunavik region of Quebec, for instance). The characters in the key above help separate the two taxa.

The typification by A. J. Richards (1985) would leave the common dandelion of both Europe and North America without a valid name (J. Kirschner and J. ?tepánek 1987). For the time being, with the nomenclatural situation still not resolved, I am following traditional usage of the name *Taraxacum officinale*.

***Taraxacum scopulorum* (A. Gray) Rydb. [FNA19, HC2]**

Mem. New York Bot. Gard. 1: 455.  
alpine dandelion

*Taraxacum lyratum* (Ledeb.) DC. [HC], Misapplied

First collected in WA in Whatcom County in 1934. Determination confirmed in 2019.

***Tetradymia* [FNA20, HC, HC2]**

Prodr. 6: 440. 1838.  
horse-brush

***Tetradymia canescens* DC. [FNA20, HC, HC2]**

Prodr. 6: 440.  
gray horsebrush, spineless horsebrush

*Tetradymia inermis* Nutt.

***Tonestus* [FNA20, HC2]**

Bot. Gaz. 37: 262. 1904.  
serpentweed

***Tonestus lyallii* (A. Gray) A. Nelson [FNA20, HC2]**

Bot. Gaz. 37: 262.  
Lyall's goldenweed, Lyall's serpentweed

*Haplopappus lyallii* A. Gray [HC]

FNA20: "Tonestus lyallii is widespread in the central Rocky Mountains and ranges of the Pacific Northwest, and is known in the Great Basin from collections in the Ruby Mountains in Elko County, Nevada. Populations documented from the Coast Range in Siskyou and Trinity counties, California, are disjunct from those in Oregon and Washington by more than 700 km."

***Townsendia* [FNA20, HC, HC2]**

Fl. Bor.-Amer. 2: 16, plate 119. 1834.  
townsend daisy, townsendia

***Townsendia florifera* (Hook.) A. Gray**

Proc. Amer. Acad. Arts xvi. 84.

showy Townsend daisy, showy townsendia

*Townsendia florifer* (Hook.) A. Gray [FNA20, HC, HC2]

*Townsendia florifer* (Hook.) A. Gray var. *florifer*

*Townsendia florifera* (Hook.) A. Gray var. *watsonii* (A. Gray) Cronquist

***Tragopogon* [FNA19, HC, HC2]**

Sp. Pl. 2: 789. 1753; Gen. Pl. ed. 5, 346. 1754.

goatsbeard, salsify

***Tragopogon dubius* Scop. [FNA19, HC, HC2]**

Fl. Carniol. ed. 2. 2: 95.

yellow salsify

FNA19: "<i>Tragopogon dubius</i>" is naturalized across much of North America. It typically grows in sites drier than those where <i>T. pratensis</i> is found."

***Tragopogon floccosus* Waldst. & Kit. [HC2]**

Descriptiones et Icones Plantarum Rariorum Hungariae 2: 116, t. 112.

woolly goatsbeard

***Tragopogon mirus* Ownbey [FNA19, HC, HC2]**

Amer. J. Bot. 37: 497.

remarkable goatsbeard

FNA19: "Tragopogon mirus is allotetraploid, formed from *T. dubius* and *T. porrifolius*. It originated (probably repeatedly) in the United States (eastern Washington, adjacent Idaho, and near Flagstaff, Arizona). F1 hybrids between *T. dubius* and *T. porrifolius* resemble *T. mirus* but are less robust, have low pollen stainability, and set few, if any, seeds. *Tragopogon mirus* does not occur in Europe, but *T. dubius* and *T. porrifolius* may occasionally hybridize there when sympatric."

***Tragopogon miscellus* Ownbey [FNA19, HC, HC2]**

Amer. J. Bot. 37: 498.

hybrid goatsbeard

FNA19: "Tragopogon mirus is allotetraploid, formed from *T. dubius* and *T. porrifolius*. It originated (probably repeatedly) in the United States (eastern Washington, adjacent Idaho, and near Flagstaff, Arizona). F1 hybrids between *T. dubius* and *T. porrifolius* resemble *T. mirus* but are less robust, have low pollen stainability, and set few, if any, seeds. *Tragopogon mirus* does not occur in Europe, but *T. dubius* and *T. porrifolius* may occasionally hybridize there when sympatric."

***Tragopogon porrifolius* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 789. 1753.

oyster plant, purple salsify

FNA19: "Tragopogon porrifolius is occasionally cultivated in Europe and naturalized across much of North America. It grows typically in sites drier than those of *T. pratensis* and in sites shadier and/or moister than those of *T. dubius*. As currently circumscribed, it may not be monophyletic, and nomenclatural changes for the populations here may be required. In North America, *T. porrifolius* hybridizes with both *T. dubius* and *T. pratensis* (= *T. ×neohybridus* Farwell, described from North America, and *T. ×mirabilis* Rouy, described from Europe)."

***Tragopogon pratensis* L. [FNA19, HC, HC2]**

Sp. Pl. 2: 789.

Jack-go-to-bed-at-noon, meadow salsify

*Tragopogon pratensis* L. ssp. *pratensis*

FNA19: "Tragopogon pratensis is naturalized across much of North America. The circumscription and infraspecific taxonomy of *T. pratensis* in Europe are debated, and the name *T. pratensis* may prove to be inaccurately assigned to the introduced populations in North America." ssp. taxonomy needs more study, using key in Stace (1997)

***Tripleurospermum* [FNA19, HC2]**

Tanacetean. 31. 1844.

mayweed

***Tripleurospermum inodorum* (L.) Sch. Bip. [FNA19, HC2]**

Tanacetean. 32.

false chamomile, false mayweed, scentless mayweed

*Matricaria inodora* L. [HC]

*Matricaria maritima* L. ssp. *inodora* (L.) Soó

*Matricaria perforata* Mérat

*Tripleurospermum maritimum* (L.) W.D.J. Koch ssp. *inodorum* (L.) Applequist

*Tripleurospermum perforatum* (Mérat) M. Lainz

FNA19: "Tripleurospermum inodorum has been classified as a noxious weed (class C) in the state of Washington and is considered invasive in other states (it is resistant to some herbicides); it is a weed of cereals in western Canada. W. L. Applequist (2002) has shown that the name *Matricaria inodora* is not a superfluous new name for *M. chamomilla* as earlier stated by S. Rauschert (1974). Therefore, the appropriate name under *Tripleurospermum* is *T. inodorum*. She also considered its type to belong in *T. maritimum* and formally recognized it there as subsp. *inodorum*, on the basis of hybridization with other *T. maritimum* subspecies (A. Vaarama 1953); on the same basis, however, Hämet-Ahti maintained the species distinction between *T. inodorum* and *T. maritimum*, while making *T. phaeocephalum* a subspecies of the latter. Q. O. N. Kay (1994), in a more extensive review of the literature and of hybridization data, also maintained *T. inodorum* and *T. maritimum* as distinct species, a conclusion followed here. From the standpoint of weed science, taxonomic merging of *T. inodorum* and *T. maritimum* has the inconvenience of grouping under a single specific name taxa that have different physiologies, ecologies, weed potentials, and, possibly, reactions to weed control measures. The name *Matricaria inodora* var. *agrestis* Weiss was not validly published."

***Tussilago* [FNA20, HC, HC2]**

Sp. Pl. 2: 865. 1753; Gen. Pl. ed. 5, 372. 1754.

coltsfoot

***Tussilago farfara* L. [FNA20, HC, HC2]**

Sp. Pl. 2: 865.

coltsfoot

FNA20: "Flowering heads of *Tussilago farfara* close at night (laminae of ray corollas arch and roll inward). The species is becoming an invasive weed in some areas."

***Uropappus* [FNA19, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 424. 1841.

silverpuffs

***Uropappus lindleyi* (DC.) Nutt. [FNA19, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 425.

Lindley's false silverpuffs

*Microseris lindleyi* (DC.) A. Gray [HC]

*Microseris linearifolia* (Nutt.) Schultz Bipontinus

*Uropappus linearifolius* Nutt.

FNA19: "Uropappus lindleyi was placed in *Microseris* (K. L. Chambers 1955) because of two allotetraploid species formed by hybridization with annual members of that genus. A number of morphologic features, including narrow, acuminate leaves with villous-ciliate margins, erect heads, relatively long outer phyllaries, cypselae often short-beaked, and pappi of white, lustrous scales suggest a connection with *Nothocalaïs*, especially *N. troximoides*. Phylogenetic studies of chloroplast DNA variation (R. K. Jansen et al. 1991b; J. Whitton et al. 1995) link *Uropappus* with *Nothocalaïs* and *Agoseris* as a sister clade to *Microseris*. Consequently, Jansen et al. separated *Uropappus* from *Microseris* and placed the two allotetraploid species in *Stebbinsoseris*."

***Wyethia* [FNA21, HC, HC2]**

J. Acad. Nat. Sci. Philadelphia. 7: 39, plate 5. 1834.

mule's-ears, wyethia

***Wyethia amplexicaulis* (Nutt.) Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 352.  
northern mule's ears, smooth dwarf sunflower

***Wyethia angustifolia* (DC.) Nutt. [FNA21, HC, HC2]**

Trans. Amer. Philos. Soc., n. s. 7: 352.  
narrowleaf mule's ears, narrowleaf wyethia

*Alarconia angustifolia* DC.

*Wyethia angustifolia* (DC.) Nutt. var. *angustifolia*

*Wyethia angustifolia* (DC.) Nutt. var. *foliosa* (Congdon) H.M. Hall

***Xanthium* [FNA21, HC, HC2]**

Sp. Pl. 2: 987. 1753; Gen. Pl. ed. 5, 424. 1754.  
cocklebur

***Xanthium orientale* L. [WTU]**

cocklebur, common cocklebur

*Xanthium strumarium* L.

*Xanthium strumarium* L. var. *canadense* (Mill.) Torr. & A. Gray

*Xanthium strumarium* L. var. *glabratum* (DC.) Cronquist

*Xanthium strumarium* L. var. *oviforme* (Wallr.) M. Peck

*Xanthium strumarium* L. var. *pensylvanicum* (Wallr.) M. Peck

*Xanthium strumarium* L. var. *strumarium*

*Xanthium strumarium* L. var. *wootonii* (Cockerell) W.C. Martin & C.R. Hutchins, invalidly published

*Xanthium strumarium* is a misapplied name referential to plants occurring in Eurasia, southeast Asia, and northern Africa. It is not known from North America..

***Xanthium spinosum* L. [FNA21, HC, HC2]**

Sp. Pl. 2: 987.  
spiny clotbur, spiny cocklebur

*Xanthium ambrosioides* Hook. & Arn.

*Xanthium spinosum* L. var. *inerme* Bel

FNA21: "Some authors have contended that *Xanthium spinosum* originated in South America and is introduced and/or naturalized everywhere else that it is found." Considered native to California in Jepson Manual (1993).

---

**Compositae (see Asteraceae)**