

Flora of the Pacific Northwest Checklist

A checklist of the Vascular Plants of the Pacific Northwest
Hosted by the University of Washington Herbarium

Family: Nymphaeaceae

6 terminal taxa (species, subspecies, and varieties).

The Flora of the Pacific Northwest Checklist aims to be a complete list of the native and naturalized vascular plants of the Pacific Northwest, with current classifications, nomenclature and synonymy.

Taxa included in the checklist:

- Native taxa whether extant, extirpated, or extinct.
- Exotic taxa that are naturalized, escaped from cultivation, or persisting wild.
- Waifs (e.g., ballast plants, escaped crop plants) and other scarcely collected exotics.
- Interspecific hybrids that are frequent or self-maintaining.
- Some unnamed taxa in the process of being described.

Family classifications follow [APG IV](#) for angiosperms, PPG I (J. Syst. Evol. 54:563-603. 2016.) for pteridophytes, and Christenhusz et al. (Phytotaxa 19:55-70. 2011.) for gymnosperms, with a few exceptions. Nomenclature and synonymy at the rank of genus and below follows the [2nd Edition of the Flora of the Pacific Northwest](#) except where superseded by new information.

Accepted names are indicated with **blue type**, synonyms with **gray type**.

Native species and infraspecies are marked with **bold-face type**.

*Non-native and introduced taxa are preceded by an asterisk.

Please note: This is a working checklist, continuously updated. Use it at your discretion.

Created from the Flora of the PNW Checklist database on June 4th, 2026 at 10:27am PT.

Available online at <https://burkeherbarium.org/pnwflora/>

Comments and questions should be addressed to the checklist administrators:

David Giblin (dgiblin@uw.edu)

Peter Zika (zikap941@gmail.com)

Suggested citation:

LASTNAME, F., OTHER PEOPLE. 2023+. Checklist of the Vascular Plants of the Pacific Northwest. University of Washington Herbarium. <https://burkeherbarium.org/pnwflora/>. Accessed Jun 4, 2026.

Dicots:

Nymphaeaceae [FNA3, HC, HC2] Water-Lily Family

Nuphar [FNA3, HC, HC2]

Fl. Graec. Prodr. 1: 361. 1809.
cow-lily, yellow water-lily

Nuphar polysepala Engelm. [FNA3, HC2]

Trans. Acad. Sci. St. Louis. 2: 282. 1865.
yellow pond lily, spatterdock

Nuphar lutea (L.) Sm., misapplied

Nuphar lutea (L.) Sm. ssp. *polysepala* (Endelm.) E.O. Beal [KZ99]

Nuphar polysepalum Engelm. [HC]

FNA3:Plants intermediate between *Nuphar polysepala* and *N . variegata* occur in eastern British Columbia."

Nuphar variegata Durand [FNA3, HC2]

Rep. (Annual) Regents Univ. State New York State Cab. Nat. Hist. 19: 73. 1866.

Nuphar variegatum Engelm. [HC], orthographic variant

Nymphaea [FNA3, HC, HC2]

Sp. Pl. 1: 510. 1753; Gen. Pl. ed. 5, 227, 1754.
water-lily

**Nymphaea alba* L. [HC2]

Nymphaea leibergii Morong [FNA3, HC2]

Bot. Gaz. 13: 124, in note. 1888.

**Nymphaea odorata* Aiton [FNA3, HC, HC2]

Hort. Kew. 2: 227. 1789.
American water-lily, fragrant water-lily

*ssp. *odorata* [FNA3, HC2]

Castalia lekophylla Small

Nymphaea odorata Aiton f. *rubra* (E. Guillon) Conard

Nymphaea odorata Aiton var. *gigantea* Tricker

Nymphaea odorata Aiton var. *godfreyi* D.B. Ward

Nymphaea odorata Aiton var. *odorata*

Nymphaea odorata Aiton var. *rosea* Pursh

Nymphaea tetragona Georgi [FNA3, HC, HC2]

Bemerk. Reise Russ. Reich. 1: 220. 1775.

pygmy water-lily

Possibly extirpated. FNA3: "Although broadly distributed in the northwest part of the flora, *Nymphaea tetragona* is apparently not common over the Canadian portion of its range. It was collected once in extreme northwestern Washington but is believed to be extirpated there. True *N . tetragona* is absent from northeastern North America and, now, from the conterminous United States, where this name has usually been applied to what is here segregated as *N . leibergii* . In size and shape of leaves and flowers the two taxa are very similar. They differ in the leaf mottling often present in developing leaves of *N . tetragona* but absent in *N . leibergii* ; the distinctly tetragonal appearance of the receptacle in *N . tetragona* ; and in the longer carpellary appendages, the presence usually of more stamens, and purple-colored stamens and pistils in *N . tetragona* . Only in living plants is it apparent that leaves of *N . leibergii* are thicker with impressed veins abaxially compared to the relatively thin leaves with raised veins in *N . tetragona* . Although distinctions in sepal and petal apices (often acute in *N . tetragona* and often rounded in *N . leibergii*) were the basis for the establishment of *Castalia leibergii* , the characters are variable in both taxa

and thus of limited utility in distinguishing them."